PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 52

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABBOTT, E. V., (449)
ADDISON, E. A., (559)
AFANASIEV, M. M., 184
AL-BELDAWI, A. S., 781
ALLEN, ROSS M., 935
ALLOW, JASSIM M., (626)
AL-SHUKRI, MAHDI MAJID, 910
ALTMAN, JACK, (908)
AMIN, K. S., (300)
AMLING, H. J., (747)
ANAND, S. C., (569)
ANDERSON, GERALD W., 328, 538
ANDERSON, J. L., (492)
ANDERSON, P. M., (678)
ANDERSON, RALPH L., (538)
ANDERSON, ROBERT L., 811
ANZALONE, LOUIS, Jr., 705, 775
ARAGAKI, M., (405), (946)
ARGUELLES-RASA, ENGRACIA, 734
ASHOUR, W. E., (79), (84)
AULAKH, KIRPAL S., 555

BABINSKI, P. L., (635)
BAGGA, H. S., 582, 835
BAJPAI, G. K., (118), (963)
BANERJEE, A. K., 118, (367)
BARDSLEY, H. C., (339)
BARKAI-GOLAN, RIVKA, (472)
BARKSDALE, T. H., 284
BARRAT, J. G., 91, (287)
BEEREBOOM, J., (479)
BELL, D. K., 240
BENIWAL, S. P. S., 817
BENZIE, J. W., (37)
BERAHA, L., 167, 974
BHARGAVA, K. S., 477
BIRCHFIELD, WRAY, 127, (423), 786
BITTERS, W. P., (952)
BLANPIED, G. D., 865, 867
BLOMSTROM, ROY, (305)
BOLING, MAX, (611)
BONINSEGNA, J., (674)
BOONE, D. M., (341)
BOOTH, J. A., (944)
BOSE, ELAINE, (722)
BOYLE, JOHN S., (972)
BRANN, J. L., (469)
BRAUN, ALVIN J., (386)
BRAVERMAN, SAMUEL W., 235
BRITTON, MICHAEL P., (922)
BRODIE, BILL B., 19, 193
BROWDER, L. E., 148
BROWN, A. G., (799)
BROWN, G. B., (486)
BROWN, G. ELDON, 844
BRUEHL, G. W., 590, (716)
BURLEIGH, J. R., (186)
BURTON, C. L., 847
BURTON, G. W., (46)
BUTCHER, CLYDE L., 295

CAFLEY, J. D., (692)
CAIRNS, ELDON J., (40)
CAMPBELL, R. N., 379
CANTWELL, ALLAN M., 957
CAPPELLINI, R. A., 479
CARROLL, R. B., (882)
CARROLL, T. W., (201)
CAVINESS, C. E., (355)
CHALKLEY, J. H., (467)
CHAND, J. N., 145
CHANDLER, W. A., 695
CHAPMAN, R. A., (204)
CHARLES, F. M., (890)
CHAUBE, H. S., (446)
CHAUHAN, L. S., 963
CHEE, K. H., 5, 132
CHENEY, P. W., (499)
CHEO, P. C., 639
CHI, C. C., 939
CHOOPANYA, DUANGCHAI, 926
CHRISTIE, S. R., 763
CLARK, FRED, (887)
CLARK, R. L., 17
COHEN, EDWARD E., (350)
COHEN, Y., (310)
COHN, E., 525
COLE, H., 410, 545, 678
COLE, JOHN R., 947
CONNOR, S. R., 654
COOK, A. A., 381
COOK, R. JAMES, 716
CORDELL, C. E., 37
COX, R. S., 339
CRAIG, JEWEUS, 134, (136)
CULP, T. W., 522, 914

DANCE, B. W., 659
DANESH, DARIUSH, (687)
DANIEL-KALIO, L. A., (134)
DARRAG, I. E. A., 399
DAVIDSON, R. W., (820)
DAVIS, J. R., (751)
DAVIS, R. T., 419
DAVIS, TERRY C., 496
DEAN, C. E., 887

PLANT DISEASE REPORTER INDEX

Issued by

CROPS RESEARCH DIVISION

Plant Industry Station
Volume 52

Beltsville, Maryland
Number 13

AUTHOR INDEX

ABBOTT, E. V., (449)
ADDISON, E. A., (559)
AFANASIEV, M. M., 184
AL-BELDAWI, A. S., 781
ALLEN, ROSS M., 935
ALLOW, JASSIM M., (626)
AL-SHUKRI, MAHDI MAJID, 910
ALTMAN, JACK, (908)
AMIN, K. S., (300)
AMLING, H. J., (747)
ANAND, S. C., (569)
ANDERSON, GERALD W., 328, 538
ANDERSON, J. L., (492)
ANDERSON, P. M., (678)
ANDERSON, RALPH L., (538)
ANDERSON, ROBERT L., 811
ANZALONE, LOUIS, Jr., 705, 775
ARAGAKI, M., (405), (946)
ARGUELLES-RASA, ENGRACIA, 734
ASHOUR, W. E., (79), (84)
AULAKH, KIRPAL S., 555

BABINSKI, P. L., (635)
BAGGA, H. S., 582, 835
BAJPAI, G. K., (118), (963)
BANERJEE, A. K., 118, (367)
BARDSLEY, H. C., (339)
BARKAI-GOLAN, RIVKA, (472)
BARKSDALE, T. H., 284
BARRAT, J. G., 91, (287)
BEEREBOOM, J., (479)
BELL, D. K., 240
BENIWAL, S. P. S., 817
BENZIE, J. W., (37)
BERAHA, L., 167, 974
BHARGAVA, K. S., 477
BIRCHFIELD, WRAY, 127, (423), 786
BITTERS, W. P., (952)
BLANPIED, G. D., 865, 867
BLOMSTROM, ROY, (305)
BOLING, MAX, (611)
BONINSEGNA, J., (674)
BOONE, D. M., (341)
BOOTH, J. A., (944)
BOSE, ELAINE, (722)
BOYLE, JOHN S., (972)
BRANN, J. L., (469)
BRAUN, ALVIN J., (386)
BRAVERMAN, SAMUEL W., 235
BRITTON, MICHAEL P., (922)
BRODIE, BILL B., 19, 193
BROWDER, L. E., 148
BROWN, A. G., (799)
BROWN, G. B., (486)
BROWN, G. ELDON, 844
BRUEHL, G. W., 590, (716)
BURLEIGH, J. R., (186)
BURTON, C. L., 847
BURTON, G. W., (46)
BUTCHER, CLYDE L., 295

CAFLEY, J. D., (692)
CAIRNS, ELDON J., (40)
CAMPBELL, R. N., 379
CANTWELL, ALLAN M., 957
CAPPELLINI, R. A., 479
CARROLL, R. B., (882)
CARROLL, T. W., (201)
CAVINESS, C. E., (355)
CHALKLEY, J. H., (467)
CHAND, J. N., 145
CHANDLER, W. A., 695
CHAPMAN, R. A., (204)
CHARLES, F. M., (890)
CHAUBE, H. S., (446)
CHAUHAN, L. S., 963
CHEE, K. H., 5, 132
CHENEY, P. W., (499)
CHEO, P. C., 639
CHI, C. C., 939
CHOOPANYA, DUANGCHAI, 926
CHRISTIE, S. R., 763
CLARK, FRED, (887)
CLARK, R. L., 17
COHEN, EDWARD E., (350)
COHEN, Y., (310)
COHN, E., 525
COLE, H., 410, 545, 678
COLE, JOHN R., 947
CONNOR, S. R., 654
COOK, A. A., 381
COOK, R. JAMES, 716
CORDELL, C. E., 37
COX, R. S., 339
CRAIG, JEWEUS, 134, (136)
CULP, T. W., 522, 914

DANCE, B. W., 659
DANESH, DARIUSH, (687)
DANIEL-KALIO, L. A., (134)
DARRAG, I. E. A., 399
DAVIDSON, R. W., (820)
DAVIS, J. R., (751)
DAVIS, R. T., 419
DAVIS, TERRY C., 496
DEAN, C. E., 887

DEAN, LESLIE L., (295), (534)
DeBLOIS, L. DONALD, (496)
DEEP, IRA W., 102
DELP, C. J., 95
DEMSKI, JAMES W., 972
deSWARDT, G. H., (479)
DeWOLFE, T. A., (281), (952)
DIAB, K. A., 149, 150
DICKENS, L. E., 225, (542)
DICKERSON, O. J., (559)
DOCAMPO, D., (703)
DODGE, F. N., (189)
DOLLINGER, E. J., (863)
DOWLER, W. M., 907
DRIVER, CHAS. H., 370
DUICH, J., (410)

EDWARDSON, J. R., (763)
EL-ERAKI, S., (149), (150)
ELLETT, C. W., (244)
EL NUR, E., 405
ENGLISH, HARLEY, (751)
EPPS, JAMES M., 45
ERCEGOVICH, C. D., (545)
ERWIN, D. C., 281
ESSER, R. P., (455), 550
ETTINGER, G., (507)
EVANS, G., 904, 976
EVERSMEYER, M. G., 186

FASSULIOTIS, GEORGE, 571
FATHI, SAAD M., (79), (84)
FELDMAN, J. M., (674), (676)
FENN, O. K., (829)
FENWICK, HARRY S., (726)
FERREE, R. J., (907)
FERRIS, JOHN M., 519
FIELDHOUSE, D. J., (683)
FILER, T. H., Jr., 170, 483
FINDLEY, W. R., (863)
FISHER, K. D., 801
FLINK, PAUL R., (538)
FLOWERS, R. A., (71), (204), (589)
FOGLE, H. W., (91), (287)
FOONG, K. M., (5)
FORDYCE, CLAUDE, Jr., 712
FRAZIER, NORMAN W., 64
FREDERIKSEN, R. A., (164)
FRENCH, D. W., (538)
FRIEND, R. J., 341
FUNK, C. R., (635)

GABRIELSON, R. L., 806
GARBER, M. J., (952)
GARUD, A. B., 436
GAY, J. DAN, 416
GERDTS, M. H., (890)
GHAFFAR, A., 223
GIBRAIL, M. A., (149)
GILL, D. L., 949
GILMER, R. M., (268)
GILMOUR, C. M., (3)
GINNS, JAMES H., Jr., 579
GINSBURG, L., (479)
GLAZE, N. C., (193)
GOHEEN, A. C., (372)
GOLDEN, A. MORGAN, 423
GOOD, J. M., (193)
GOVINDARAJAN, T. S., (773)
GRABE, D. F., (459)
GRACIA, OLGA, 595, 674, (676)
GRANEK, IRVING, 479, 827
GREATHEAD, A. S., (890)
GREWAL, J. S., (300)
GRIFFIN, G. D., 492, 956
GROGAN, C. O., 611
GROVER, RAJENDRA K., (555), (602)
GUDAUSKAS, ROBERT T., (34), (509)
GUIDROZ, GERALD, (780)
GUTTER, Y., 777

HAGEDORN, D. J., 160, (190)
HAGLUND, WILLIAM A., 72, (806)
HALISKY, PHILIP M., 496, 635
HAMMETT, K. R. W., 754
HARDING, PAUL R., Jr., 245, 433, 623
HARDISON, JOHN R., 205
HARMON, S. A., (108)
HARPER, F. R., 565
HARRISON, ARTHUR N., 34, 509
HART, W. H., (890)
HAWKSWORTH, FRANK G., 125, 804
HEALD, C. M., 46
HELTON, A. W., 154
HENDRIX, FLOYD F., Jr., 334, (577)
HEUBERGER, J. W., (654)
HEWETSON, F. N., (292)
HILDEBRAND, E. M., (475), 967
HILDEBRANDT, A. C., (600)
HINDS, T. E., 820
HIRATSUKA, Y., 650
HOCKING, D., 628
HOFFMAN, J. C., (486)
HOFFMANN, J. A., 569
HOITINK, HARRY A. J., (615)
HOLCOMB, G. E., 120
HOLTZMANN, OLIVER V., 56, (405), 515
HOPPER, W. E. R., (169)
HORST, R. K., 615, 840
HOSEIN, I., 741
HSI, DAVID C. H., 365
HUBER, DON, (716)
HUNTER, R. E., 14, 158
HUTCHISON, JAMES A., 395
HYRE, R. A., 507

IBRAHIM, G., (405)

JAWORSKI, C. A., (193), (330)
JENKINS, ANNA E., 277
JENSEN, HAROLD J., 3, 169
JERATH, M. L., 237

JOFFE, A. Z., 608, 718
JOHN, V. T., 871
JOHNSON, ALVA W., 109, 114
JOHNSON, FRED A., (339)
JONES, JOHN P., 931
JONES, L. S., (924)
JONES, R. K., (907)
JORGENSON, E. C., (492)
JOSHI, M. M., (446)

KAISER, WALTER J., 687
KALOOSTIAN, G. H., 363, 924
KAMAL, (477)
KANANEN, D. L., (553)
KATCHO, ZUHAIR A., 626
KATSANOS, R. A., 68
KEIL, HARRY L., 259
KENDRICK, E. L., (569)
KenKNIGHT, G. E., (189), 307
KENNETH, R. G., 472
KEPLINGER, J. A., 386
KHERA, S., (396)
KILPATRICK, R. A., 209
KIM, S. H., (683), (685)
KINGSLAND, GRAYDON C., 430, 792
KIRKPATRICK, HUGH C., 499
KLARMAN, W. L., 792
KLINE, D. M., 605, (879)
KLOPPING, H. L., (95)
KLOTZ, L. J., 952
KNORR, L. C., (31)
KOCHAN, W. J., (154)
KOENIGS, JEROME W., 597
KOIKE, H., 646
KOZELNICKY, GEORGE M., (320)
KREBILL, R. G., 336
KRIKUN, J., (525)
KUHN, C. W., 320, 467
KUSHMAN, L. J., 475

LAFERRIERE, L., (295)
LALLMAHOMED, G. M., 136
LAMBERTI, FRANCO, 748
LAMEY, H. A., (775)
LANGDON, K. R., (550)
LASTRA, R., 171
LATHAM, A. J., 327, 747
LAUGHLIN, C. W., (162)
LAWRENCE, F. J., 811
LEAPHART, CHARLES D., 6
LERT, PETER J., (415)
LEWIS, F. H., 292
LINDNER, R. C., (499)
LINDQUIST, C. H., (829)
LISKER, N., (718)
LITTRELL, R. H., (179)
LOCK, W., (961)
LOCKE, SETH BARTON, 272, 277
LOMBARD, FRANCES F., (820)
LORBEER, J. W., (469), (581)
LORING, L. B., (169)
LOUIE, RAYMOND, 581
LOWNSBERY, B. F., 890
LUKEZIC, F. L., (882)

MacKENZIE, D. R., (24), 28, (545)
Mac NISH, G. C., 280
MacSWAN, IAIN C., (102)
MAHADEVAN, A., (703)
MALEK, R. B., 795
MALOY, OTIS C., 489, 983
MANJI, B. T., (722)
MANKAU, R., 315, 851
MARSH, PAUL B., 671
MARTIN, GEORGE C., 99
MARTIN, W. J., (127)
MARTINEZ, A. P., (550)
MARUYAMA, P. J., (650)
MARX, D. H., (577)
MASSIE, L. B., (410)
MATTHEE, F. N., (479)
McCAIN, A. H., (415)
McCARTER, STATES M., 179, 330, (416)
McCARTNEY, WILLIAM O., 198
McCLURE, MICHAEL A., 575
McCRACKEN, F. I., (170)
McCRUM, R. C., 794
McFARLANE, J. S., 297
McKINNEY, H. H., 919
McLEAN, D. M., 486
McNEAL, F. H., 201
McNEILAN, RAY A., (102)
MEREDITH, D. S., 405
METZER, R. B., (459)
MICHEL, L. J., 205
MILLER, C. R., 984
MILLER, DOUGLAS R., 305, 391, 393, 486
MILLER, J. W., 739
MILLER, P. M., 57, 441, 728, 730, 768
MILLER, P. R., (707)
MILLER, P. W., (811)
MILLER, R. E., 553
MIRCETICH, S. M., (91), 287, (790)
MISRA, S. R., (626)
MITCHELL, D. M., (790)
MITCHELL, J. T., (890)
MOHAMED, HOSNI A., 79, 84
MOLLER, W. J., 751
MONIZ, L., (784)
MORTON, D. J., 683, 685
MOSEMAN, J. G., 463
MOSSAHEBI, HOSSEIN, (687)
MOTSINGER, RALPH E., (120)
MUKA, A. A., (469)
MUKHOPADHYAY, A. N., (715)
MULLIN, R. S., (984)

NAGAI, HIROSHI, 928
NAGEL, C. M., (349)
NAPHADE, S. R., 377
NARIANI, T. K., 595, 834

NEELY, DAN, 223
NELSON, PAUL E., (840)
NELSON, R. R., (427), (605), 879
NETZER, D., (472)
NEWELL, W. R., (455)
NIEHAUS, M. H., 633
NOME, S. F., 703

O'BANNON, J. H., (218)
OBRERO, F. P., 814, 946
OGAWA, J. M., (76), 722
OKHOVAT, MAHMOUD, (687)
OLOFSSON, JAN, 264
OSHIMA, NAGAYOSHI, (225), 542
OU, S. H., (122)

PACUMBABA, R. P., 559
PAPPELIS, A. J., (68)
PARKER, M. C., 534
PARMETER, JOHN R., Jr., 452, 707
PARRIS, G. K., 744
PARTRIDGE, A. D., 46
PATIL, L. K., 784
PETERSON, GLENN W., 357, 359, 361
PHILLIPS, D. V., 895
PICARELLI, C. J., (801)
PILLAI, S. K., 600
PINCKARD, J. A., 780, (781)
PINE, T. S., 61, 250, (253)
PLATT, W. D., (452)
PONTIS, R. E., (674), 676
POOL, R. A. F., 818
POVISH, W. R., (248)
POWELL, J. M., 409
POWELL, WILLIAM M., (109), (334), 577
POWELSON, R. L., (716)
PRATT, CHARLOTTE, 268
PRESCOTT, J. M., 437
PROVVIDENTI, R., (386), (630)
PUNTER, D., 692
PURCIFULL, D. E., 759
PURNASIRI, APATHAI, (865), (867)
PURUSHOTHAMAN, D., 703

RAABE, ROBERT D., 415
RAGUNATHAN, V., (703), 705
RAI, J. N., 175
RAI, P. V., (502)
RANDALL, HUGH, 349
RAO, A. APPA, (494)
RAU, G. J., (571)
RAWLINS, T. E., 944
RAYCHAUDHURI, S. P., (834)
REBOIS, R. V., 40
REDDY, D. V. R., (494)
REDDY, M. SUGUNAKAR, 494
REDFERN, D. B., 638
REITZ, H. J., 31
REYES, LUCAS, (164)
REYNOLDS, G., 961
RHOADES, H. L., 383, 573
RICH, SAUL, (728), (730), 732
RICHARDSON, D. L., (901)
RIFFLE, JERRY W., 52
RITTER, R. M., (863)
RIVERA, C. T., 122
ROBERTS, B. J., (512), (897)
ROCHEFORT, K. J., (87)
RODRIQUEZ, ENRIQUE, 164
ROELFS, A. P., (512)
ROMIG, R. W., 512, (642)
ROSENOW, D. T., (164)
ROSS, ELDON W., (105), 635
ROSS, J. P., 344, (895)
ROTEM, J., 310
ROTH, J. N., 215
ROUNSVILLE, G. B., (87)
ROWELL, J. B., 856
RUEHLE, JOHN L., 523, 837

SABAR, Z., 860
SAKSENA, H. K., (663)
SALAMA, EL-SAYED A., 11
SAXENA, S. C., (715)
SAXENA, V. C., (175)
SCHAFER, JOHN F., (215), (395)
SCHALLER, C. C., 189
SCHEIFELE, G. L., 424, 427
SCHENCK, N. C., 979
SCHIEBER, E., 115
SCHMITTHENNER, A. F., (244)
SCHNATHORST, W. C., 981
SCHROEDER, W. T., 630
SHALLA, T. A., (860)
SHANNON, EMROY L., (365)
SHARMA, B. C., (834)
SHARP, E. L., (818)
SHEPARD, J. F., 139
SHERIDAN, J. E., 142
SHOEMAKER, P. B., 469
SILL, W. H., Jr., (11), (559)
SILVERBORG, S. B., 105
SIMONS, M. D., 59, (205)
SIMPSON, MARION E., (671)
SIMPSON, W. R., 726, 965
SINCLAIR, J. B., (399), 667
SINGH, KISHAN, 626
SINGH, R. N., 662, 663
SINGH, R. S., 446, 602
SINHA, S. K., 715
SIRADHANA, BABU SINGH, 244
SIRRY, A. R., (79), (84)
SITTERLY, W. R., 49
SIVAPALAN, P., 528
SKELLY, J. M., 882
SKILLING, D. D., (37)
SLATTERY, ROBERT J., 692
SLEETH, BAILEY, 232
SMART, GROVER C., Jr., 455
SMERLIS, E., 167, 403, 738
SMILEY, J. H., 71, 204, 589
SMITH, F. F., (530)

SMITH, F. H., (571)
SMITH, PAUL G., (928)
SNOW, G. A., (419)
SONODA, R. M., 76
SOWELL, GROVER, Jr., (467)
SPEARS, JOSEPH F., 184
SPENCER, JAMES A., 751
SPIEGEL, SARAH, (310)
SPOTTS, ROBERT A., (908)
SRIVASTAVA, S. K., (817)
STALEY, JOHN M., (804), 908
STEINER, P. W., (469)
STEWART, D. M., 642, 897
STEWART, JAMES L., (357)
STEWART, JOSEPH K., (262)
STOKER, G. L., (956)
STOKES, D. E., 118
STOUFFER, R. F., (292)
STOVER, R. H., 901
STRANDBERG, JAMES O., 859
STROBEL, GARY A., 502
STUBBS, H. F., (467)
SUBRAMANIAN, S., 773
ŠUTIĆ, D., (250), 253

TACHIBANA, H., 459, 613
TANDON, I. N., (118), 367
TANTERE, D. M., (122)
TARTÉ, RODRIGO, 587
TAYLOR, A. L., 218
TAYLOR, G. S., (441)
TELIZ, DANIEL, 372
TEMP, M. W., 190
TEWARI, J. P., (175)
THOMAS, CLAUDE E., 374, 375, 761
THOMPSON, L. S., 213, 393
THYR, B. D., 741
TIKYANI, M. G., 396
TIPPETT, R. L., 449
TOMLINSON, HARLEY, (732)
TOMPKINS, D. R., (806)
TOOLE, E. R., (170), 482, (483), 822
TRAYLOR, JACK, (618)
TROUTMAN, J. L., (522), (914)
TRUJILLO, E. E., (814), (946)
TU, J. C., 323
TYAGI, P. D., 569

ULLSTRUP, A. J., 922

URIE, A. L., (876)

van der ZWET, T., 698
Van SICKLE, G. A., 455, (638)
VEIRS, EVAN I., (393)
VERMA, P. S., (145)
VERMA, S. C., (963)
VERRALL, A. F., 248

WAGNON, H. KEITH, 618
WALTERS, H. J., 355
WALTON, G. S., 824
WATERWORTH, H. E., 248, 530
WAY, R. D., (268)
WEBB, R. E., (761)
WEINER, ADAM C., (618)
WELLS, JOHN M., 262
WERNHAM, C. C., 24, (28), (424), (427)
WESTER, HORACE V., 350
WESTER, R. E., 563
WHITTY, E. B., (984)
WICKER, ED F., (6)
WIGGS, D. N., 528
WIHRHEIM, S. E., (57), (441)
WILLIAMS, A. S., 162
WILLIAMS, E. B., 799
WILLIAMS, F. J., 300
WILLIAMS, HAROLD E., (618)
WILLIAMS, L. E., 863
WILLIS, C. B., (213)
WILLIS, WILLIAM G., (357), (652)
WILSON, CHARLES L., 79
WILSON, E. E., 227
WILSON, ROY A., (259)
WOOD, L. S., (349)
WOOD, R. E., (370)
WORLEY, RAY E., 108
WYSONG, DAVID S., (359), (361), 652

YANG, CHARLES Y., (71)
YISRAELI, U., (525)
YOUNG, H. C., Jr., (437)

ZALASKY, H., 829
ZENTMYER, G. A., (115), (223), 790
ZETTLER, F. W., (763)
ZILLER, W. G., (336)
ZIMMER, D. E., 876
ZUCKERMAN, B. M., 87

SUBJECT INDEX

Abies spp.: rust fungus Hyalopsora aspidiotus on fir in the West, first Idaho record on fir, 336
--- concolor (white fir): dwarf mistletoe infection at girdle and segment regions of fir branches, 452; Furcaspora pinicola found on needles of white fir, a new host, 486
--- grandis (grand fir): decay fungi associated with young grand fir trees, 489
Agar (see Cultures)
Agaricus bisporus (see Mushroom)
Agrobacterium tumefaciens: 102; 492; 550
Agropyron (see Grasses)
Air pollution: petunia and Datura suffered injury due to some kind of air pollution

in Connecticut, 732; exhaust fume damage to redbud, 744; ozone damage to apple, 730; ozone damage to Pinus ponderosa in southern California, 707
Alabama: 34; 496; 747
Alberta: 565
Alfalfa: anthracnose, caused by Colletotrichum trifolii, no previous record on alfalfa from the Pacific Northwest, 983; Fusarium wilt and root rot development in relation to nutrition stress, 939
Alternaria sp.: 215
--- spp.: 248
--- longipes: 71
--- tenuis: 582
--- tenuissima: 145
Amaranthus macrocarpus: dwarf amaranth new host for Verticillium nigrescens, 904
Amendments: petroleum mulch, helped reduce damping-off of cucumber when combined with band treatments of fungicides, 806; pine sawdust, paper and cottonseed meal plus nitrogen suppressed root invasion of eggplant by Heterodera tabacum, 441; steer and alfalfa green manure in outdoor microplots reduced root-knot disease in okra, 315
Anatomical study: of effects of cadang-cadang disease on coconuts, 734
Announcements: 256; 984
Anthracnose: alfalfa, in Washington, 983
---: of pear, historical record of pear spotted anthracnose (Elsinoë piri), 277
---: of pineapple in India, first report of disease, 436
---: of sorghum, sorghum lines showed marked difference in reaction to artificial inoculation with Colletotrichum graminicola, 164
---: of tomato, yield increases and disease reduction with experimental fungicides, 685
Antibiotics: cycloheximide (Acti-dione) effective in inhibiting growth of Armillaria mellea on potato-dextrose agar, 639
---: phytoactin and cycloheximide not effective for control of blister rust disease of western white pine, 6; phytoactin, cycloheximide, methyl hydrazone, and semicarbazone not effective against eastern gall rust, did delay sweetfern rust symptom development on jack pine, 538
---: streptomycin spray controlled fire blight on apples with no detectable residue on fruit, 259
Antiserum (see Serum)
Aphanomyces euteiches: 190; 264
Aphelenchoides sp.: 52
--- besseyi: 56
--- fragariae: 118; 455
Aphid: green peach aphid, Myzus persicae, implicated as vector of russet crack virus of sweetpotato, 967
Apple: European canker, control of by eradicative and protective fungicides in California, 227
---: fire blight: control with streptomycin sprays, 259; its distribution throughout the world and its recent spread, 698
---: ozone damage to, 730
---: Penicillium rot controlled with thiabendazole as postharvest fungicide, 867; Penicillium and Botrytis rot of McIntosh apples handled in water, 865
---: scab: late season application of Fungicide 1991 and its analog completely inhibited perithecial development, 654; new physiologic race of Venturia inaequalis in England, 799; reduction of spring discharge of Venturia inaequalis ascospores by composting overwintering leaves, 728
---: streptomycin applied as spray controlled fire blight with no detectable residue on fruit, 259
---: virus, mosaic virus, Prunus virginiana a natural host for, 794
Apricot (see Prunus)
Aquatic plants: several found hosts of Aphelenchoides fragariae in Miami, Florida, thought to be first report of this nematode parasitizing aquatic plants, 455
Arachis hypogaea (see also Peanut): relative susceptibility of some peanut varieties to Cercospora personata in Sudan, 405
Arceuthobium americanum: 125, 409
--- campylopodum: 452
Argentina: 595; 674; 676
Arizona: 804
Arkansas: 79; 170
Armillaria mellea: 415; 639
Ash (see also Fraxinus): dieback, of white ash, disease development in New York, 1962-1968, 105
Aspen (see Populus)
Aspergillus spp.: 582
--- flavus: 671; 718
Australia: 280; 976
Azalea: Cylindrocladium blights controlled with drench of Fungicide 1991, 615

Bacteria: associated with wetwood of cottonwood in Mississippi, 822; Erwinia

aroideae, one of several bacteria isolated from cotton plants which cause boll rot, 14; preparation of a semi-solid agar medium for detection of pectolytic enzyme activity in Erwinia carotovora, 167; Streptomyces phage can be transmitted by saprozoic nematodes, 3
Bacterial blight (see also Blight)
Bacterial diseases: bacterial blight: of cotton, caused by Xanthomonas malvacearum, 739; 981; fire blight of apple, caused by Erwinia amylovora, 259; 698; halo blight of bean, caused by Pseudomonas phaseolicola, 225; 295; 534; of rice, caused by Xanthomonas oryzae, 663; of sugarbeet, caused by Pseudomonas aptata, 715
---: bacterial canker: citrus canker caused by Xanthomonas citri eradicated from Louisiana, 667; of tomato, caused by Corynebacterium michiganense, in Argentina, 595; of tomato, level of inoculum needed for infection, 741
---: bacterial crown gall: disease on mazzard cherry, control tests with soil fumigants, 102; crown gall of raspberry, caused by Agrobacterium tumefaciens, 492
---: bacterial leaf scald of sugarcane, caused by Xanthomonas albilineans, 646
---: bacterial rind necrosis, of watermelon in south Texas caused by Erwinia sp., 375
---: bacterial rot (see also Rot): boll rot of cotton, caused by Erwinia aroideae, 14; potato ring rot, caused by Corynebacterium sepedonicum, 502
---: bacterial spot: of pepper caused by Xanthomonas vesicatoria, effect of bactericides, saccharin and alar on, 683; of tomato, controlled successfully in Florida with proper liaison between expert and grower, 339
---: bacterial wilt: of 'Bluggoe' (cooking banana), caused by Pseudomonas solanacearum, 901; of tomato, caused by Pseudomonas solanacearum, 330
Baldcypress: Stereum taxodii causes peck in baldcypress, 482
Banana: Deightoniella torulosa, causes fruit spotting, first published record of disease in Hawaii, 405
Barberry: stem rust of wheat and oats on, in 1965, 512; 1966, 897; 1967, 642
Barley: loose smut, germination reduced during storage after anaerobic treatment, 792
---: net blotch, barley seedlings showed resistance to Pyrenophora teres, 215
---: powdery mildew, reactions of barley to Erysiphe graminis f. sp. hordei from North America, England, Ireland and Japan, 463
---: Pyrenophora-Cochliobolus (Helminthosporium) disease complex in winter barley in Indiana, 395
---: Pythium species tested for pathogenicity to oat, barley, and wheat seedlings, 209
---: root-knot nematode, Meloidogyne naasi caused stunting of barley plants in western Oregon, 169
Bean: diseases of in Colorado in 1967, 542
---: downy mildew of lima bean, two new lines G 1 and G 2 completely immune to strains A and B of Phytophthora phaseoli in 1967 tests, 563
---: halo blight: copper compounds as protective sprays beneficial in reducing incidence of halo blight in Colorado, 225; control in Idaho by use of field and windrow inspections, 295; detection of infected seed with use of ultraviolet light, 534
---: ozone damage, staining technique to differentiate dead cells injured by ozone, 732
---: Rhizoctonia solani root rot resistance in snapbean, 12 breeding lines and PI accessions showed resistance in greenhouse studies in South Carolina, 486
---: stem rot of snapbean caused by Pythium myriotylum in Georgia, apparently first report of infection under field conditions, 416
Beet (tablebeet): infection by Rhizoctonia sp. in Colorado in 1967, 542
Belonolaimus longicaudatus: 19; 46; 523
Bibliography: "Bibliography of Soybean Diseases" now available, 256
Bidens pilosa: host of Bidens mottle virus, 763
Bifusella linearis: 305
Bipolaris tetramera: 377
Black rot: of sweetpotato, control of by hot-water sprout treatment seems promising, 475
Blast disease: of rice, caused by Piricularia oryzae, new medium to induce sporulation, 703
Bleeding necrosis: of sweet gum in Illinois and Indiana, pathogenicity of Botryosphaeria ribis confirmed, 223
Blight: bacterial: of cotton, widespread increase in virulence in Missouri, several races identified, 739; introduction into California on acid-delinted and fumigated cotton seed, 981; of rice (paddy), caused by Xanthomonas

oryzae in India, 663; of sugarbeet in India, 715
---: Cercospora blight of junipers, in Nebraska, Bordeaux mixture better of two fungicides for control, 361
---: Diplodia tip blight of pines in Kansas and Nebraska, 359
---: early, of tomato, Colletotrichum phomoides incidence reduced with fungicide treatments, 685
---: fire: on apples, control during post-bloom period with streptomycin sprays, 259; of pear, apple and other rosaceous hosts, recent spread and present distribution in the world, 698
---: of grasses: 142 isolates of Cochliobolus carbonum from diverse source hosts and geographic origins tested for capacities to induce lesion or blight reactions on 28 gramineous test species, 605; Cochliobolus heterostrophus capable of inducing blight on 34 gramineous test species, 879
---: gummy stem, new symptom of on cucumber fruit, 49
---: halo: of beans, best control is production of bean seed in semi-arid West, with field and windrow inspections, 295; foliage sprays with copper compounds for prevention of spread of halo blight gave protection, 225; use of ultraviolet light as sampling aid to detect seed infected with Pseudomonas phaseolicola, 534
---: late, or leaf spot of celery, effect of weather, duration of high humidity, and spore load upon infection, 142
---: leaf: of Crotalaria juncea from India, 377; of papaya, a new record of disease of papaya caused by Helminthosporium rostratum, 784; leaf or seedling of corn, chlorotic lesion resistance to Helminthosporium maydis found in corn selections in Nigeria, 134
---: seedling: of corn, factors affecting severity of in United Arab Republic, 79; of corn, Nigrospora oryzae, Penicillium graminearum and two Fusarium species found pathogenic and to cause seed rot, seedling blight or both, in United Arab Republic, 84; of cotton, Fungicide 1991 effective as control fungicide, 781
---: sheath of rice, seven strains in Taiwan similar but differing in pathogenicity, 323
---: southern corn leaf blight, corn ears shipped from Florida infected, 847
Blue mold: of tobacco: in Kentucky in 1967, 204; plants exposed to low soil moisture seemed more susceptible to infection, 310
'Bluggoe' (cooking banana): Fusarium oxysporum cubense race 2 and Pseudomonas solanacearum extremely pathogenic to, Philippine accession 'Pelipita' found to replace it, 901
Boron: toxicity in greenhouse-grown chrysanthemums, 824
Botryosphaeria berengeriana: 947
--- ribis: 223
Botrytis sp.: 479; 865
--- allii: 974
--- cinerea: 248
--- squamosa: 469
Bovine: Meloidogyne arenaria survived passage through digestive tract of Jersey cow, 99
British Columbia: 579
Brown rot: of peach, Fungicide 1991 outstanding in 1967 as preharvest spray for brown rot control, 695
Brown spot: of tobacco, caused by Alternaria longipes, first reported occurrence in Kentucky, 1966, 71
Buckeye: California, apparently new record as host of European mistletoe in California, 198
Buckeye rot (see Rot)
Bunch disease: of pecan, of unknown cause, attempts to spread by inoculation ineffective, 307
Bunt: of wheat: new pathogenic race of Tilletia foetida found in Idaho, designated L-16, 569
Butt rot: decay of Engelmann spruce in Colorado due to Lentinellus montanus, 820

Cabbage (see Crucifers)
Cacao (Theobroma cacao): pod rot or black pod caused by Phytophthora palmivora, stem and root inoculation methods for testing for resistance described, 790
Cactus: cactus cyst nematode found on Echinocereus pentalophus and Echinopsis multiplex in Georgia for first time, 114
Cadang-cadang disease: of coconut, anatomic effects of, in Philippines, 734
California: 76; 198; 223; 262; 297; 305; 391; 393; 452; 618; 707; 722; 751; 860; 928; 981
Callus cultures: geranium plants successfully grown from apical meristem and callus cultures, 600
Camellia sinensis: decline in young tea fields caused by nematode, Radopholus similis, in Ceylon, 528
Canada: 409; 638; 659; 829

Candida krusei: 433
Canker: bacterial: of citrus, eradication program in Louisiana, 667; of tomato, 595; 741
---: Coniothyrium canker found on ornamental pear (Pyrus calleryana), 792
---: Cytospora, on 'Italian' prune, preinfection treatment of young trees with systemic-chemical paints, 154
---: European, of apple in California, eradicative and protective fungicides tested for control, 227
---: Populus: Hypoxylon pruinatum induced cankers on trembling aspen in Ontario, 659; Septoria musiva incidence high in 12 clones of naturally infected poplars in Canada, 829
---: Scleroderris canker, of pine, red and jack pines more susceptible than Scotch pines in outplanting study in Michigan, 37
---: stem and pink stain of teak in Tanzania associated with Fusarium solani, 628
Cantaloup (see Cucurbits)
Capsicum spp. (see also Pepper): virus disease resistance in some species from South America, 381
Carnation: Armillaria stem rot, infection of carnations planted in former prune orchard, believed to be first report of fungus on carnation, 415; Fusarium stem rot, losses from in commercial production, 840
Carrot: diseases of in Colorado in 1967, 542
Cedar-apple rust: control studies with fungicides on Juniperus scopulorum and J. virginianae, in Illinois, 692
Celery: late blight or leaf spot, conditions for infection of celery by Septoria apicola, 142
Cephalosporium sp.: 170
--- diospyri: 79
--- gramineum: 590; 818
--- gregatum: 895
Ceratocystis fagacearum: 357; 811
--- fimbriata: 475
--- ulmi: 46
Cercospora kikuchii: 931
--- personata: 405
--- sequoiae: 361
Ceylon: 528
Chemicals: mixtures of specific pesticides tested with broad spectrum soil fumigants on tomatoes, 193
---: boron toxicity to chrysanthemums in Connecticut, 824
---: fixed copper (cupric hydroxide), tribasic copper sulfate, saccharin and alar showed no significant differences for bacterial spot control on peppers, 683
---: cycloheximide, 2,4-dichlorophenoxyacetonitrile, and acrizane inhibited growth of Armillaria mellea on potato-dextrose agar, DMSO in agar showed some inhibitory effect, 639
---: 2,4-DB (4(2,4-dichlorophenoxy)butyric acid) Phytophthora resistant soybean varieties showed resistance to damage from herbicide, 355
---: DBCP nematocide in combination with different fungicides decreased Pythium population in the control of feeder root necrosis of pecans, 577
---: DMSO (dimethyl sulfoxide) can affect number of local lesions induced by tobacco mosaic virus, 61
---: phorate most effective of 11 insecticides for control of phyllody and leaf curl of Sesamum orientale in India, 367
---: salt (sodium chloride) damage to vegetation after winter-salting of roadways in Washington, D. C. area, 350
Chemurgic crops: Hibiscus, Vernonia, Tephrosia and Limnanthes found susceptible to common plant viruses, 530
Chenopodium album: new host of Verticillium nigrescens, 904
Cherry (see also Prunus): crown gall, of mazzard cherry in Oregon, soil fumigant Trizone appeared to give some control, 102
--- virus: cherry rasp-leaf virus, investigations of in California, 618; crinkle, occurrence of normal foliage in irradiated sweet cherries with crinkle, 268; necrotic ringspot virus in Pennsylvania orchard, 972
Chile: 703
Chrysanthemum: occurrence of boron toxicity, 824; Sclerotinia flower rot, an unusual occurrence in Louisiana, 120
--- viscidi-hirtum: plant introduction in SIPE program susceptible to Alternaria tenuis, 248
Circulifer tenellus: 17
Citrus: canker, caused by Xanthomonas citri, successful eradication program in Louisiana, 667
---: crinkle-scurf, found to cause reduced growth, fruit size, juice content and crop yield of Valencia orange, 31
---: gummosis, some factors in resistance of citrus to Phytophthora spp., 952
---: lemon, Candida krusei as primary pathogen, first published record of such a relationship, 433

---: postharvest decay: Penicillium mold decay on oranges and lemons as affected by ozone in storage rooms, 245; comparison of three fungicides for control of Penicillium molds of citrus fruits, 623; Penicillium digitatum decay of Valencia oranges controlled with 5-aceto-8-hydroxyquinoline sulfate + 1.0% wetting agent, 777; Fungicide 1991 outstanding as preharvest fungicide for control of postharvest decay, by Penicillium spp. and Diplodia natalensis, 844
---: virus: exocortis virus in Ortanique orange in Jamaica, found for first time, 741; reported for first time from India, 834; may be readily transmitted by contaminated knife blades, 935
Cladosporium effusum: 307
--- herbarum: 248
Cleonia lusitanica: recent plant introduction infected with Botrytis cinerea, 248
Clipping: of tomato plants, greenhouse study indicates bacterial wilt of tomato can spread from diseased to healthy plants by clipping, 330
Clover: red, root decay, more decay developed when clover root curculios were present with Fusarium spp. than with Fusarium alone, 213
Clusterbeans (Cyamopsis tetragonoloba): incidence of Alternaria leaf spot apparently correlated with temperature and rainfall in India, 145
Cochliobolus carbonum: 605
--- heterostrophus: 847; 879
--- sativus: little evidence of seedling leaf resistance in barley, 215; 395
Coconut: cadang-cadang disease, phloem necrosis associated with, 734
Coffee: Myrothecium stem necrosis and leaf spot, in Guatemala, 115; new root disease caused by Cylindrocarpon tenue, 773
Colletotrichum sp.: 436
--- dematium f. truncata: found on alfalfa near Moses Lake, Washington, new report, 983
--- graminicola: 68; 164
--- higginsianum: 248
--- phomoides: 685
--- trifolii: new record of Colletotrichum on alfalfa from Pacific Northwest, 983
Colorado: 125; 542; 820; 908
Commercial implementation: of disease control, effective in Florida, 339
Coniothyrium sp.: 792
Connecticut: 728; 730; 732; 801; 824
Corn: chlorotic lesion, resistance found in corn inbreds in Nigeria, 134
---: Diplodia stalk rot, inoculation of plants for reaction to Diplodia stalk rot with CO_2-powered BB pistol, 611
---: downy mildew, more evidence on the seedborne nature of corn downy mildews, 446
---: head smut (Sphacelotheca reiliana), controlled in Idaho with in-furrow application of 1-chloro-2-nitropropane (NIA 5961) + PCNB, 726
---: root rot caused by Helminthosporium pedicellatum, first report of occurrence in India, 477
---: rust, in Nigeria two physiologic races of Puccinia polysora found, 136
---: seedling blight: Fusarium moniliforme, F. graminearum, Nigrospora oryzae and Penicillium oxalicum found to cause seed rot, seedling blight or both, in the United Arab Republic, 84; in United Arab Republic, factors affecting severity of blight, 79
---: smut (Ustilago maydis), incidence apparently related to low temperatures, 922
---: southern corn leaf blight, on sweet corn ears shipped from Florida, first report of, 847
---: Trichometasphaeria turcica (Helminthosporium turcicum) survival capacity of simple and complex races in mixtures for several generations on a simple inbred line of corn with no genes for vertical resistance, 427
---: virus: maize dwarf mosaic virus: air-blast inoculation of corn varieties with MDMV, field studies in Pennsylvania, 24; air-blast inoculation of corn seedlings with MDMV, effect of buffer, abrasive, and dilution of inoculum on infection, greenhouse studies, 28; MDMV identified in Georgia and found in seven counties, 320; MDMV in New Mexico, first occurrence, 365; resistance of dent corn synthetics to mosaic viruses and other maize diseases, 424; two strains of MDMV recovered from dent and sweet corn in New Jersey, 496; virus-host-soil pesticides study, disulfoton, zinc + maneb, atrazine and propazine used as drenches increased grain yield in Pennsylvania, 545; virus seed transmission studies, 863
Corrections: 416; 664
Corynebacterium michiganense: 595; 741
--- sepedonicum: 502
Cotton: reduction in yield caused by diseases in 1967, 314
---: bacterial blight: acid-delinted and

and fumigated cotton seed not free of Xanthomonas malvacearum, 981; race buildup of Xanthomonas malvacearum in southeast Missouri, 739
---: boll rot: bacterium Erwinia aroideae caused severe boll rot, 14; geographic distribution in 1967 cotton crop of fibers with fluorescent spots, caused by Aspergillus flavus, 671; Phytophthora boll rot, first known record of disease in Louisiana, 780; relative frequency of different fungi with boll rot in Mississippi Delta, Fusaria, Alternaria tenuis, Aspergilli, and Diplodia gossypina most prevalent, 582; systemic fungicides and insecticides tested by quick laboratory method for control of boll rot, 835
---: Fusarium oxysporum f. vasinfectum and Verticillium dahliae interaction on Gossypium hirsutum var. skalleridis, 910
---: nematodes: Columbia lance nematode pathogenic to cotton in South Carolina, 571; reniform nematode control in Louisiana by nematocides, 786
---: root rot, caused by Phymatotrichum omnivorum, DMTT and SMDC best of 10 chemicals for penetrating soil and killing hyphae, 232
---: seedling blight caused by Rhizoctonia solani controlled with Fungicide 1991, 781; chemotherapeutic activity of certain fungicides against Rhizoctonia solani in seedlings tested, 399
---: sooty mold on cotton and other hosts, caused by Microxyphiella hibiscifolia in India, 602
---: Verticillium wilt: control effective with use of Tween 20 as additive to systemic fungicides, 944; lack of control with three systemic insecticides, 281
Cottonwood: wetwood, rod-shaped, gram-negative bacterium consistently isolated from affected trees, 822
Cotyledon necrosis: of soybean, in Iowa, believed to be noninfectious disease due to weathering, 459
Cover crop: Crotalaria spectabilis and Sesbania exaltata as summer cover crops, 573
Cowpea: identification of viruses isolated from cowpeas in Alabama, 34; in greenhouse tests Clay cultivar showed tolerance to three viruses known to infect cowpeas, 509
Crambe juncea: Cladosporium fulvum and Alternaria brassicae pathogenic to new plant introduction, 248
Cranberry: control of fairy ring disease obtained with ferbam applied immediately after harvest, 87; dieback, Diaporthe vaccinii associated with in Wisconsin, 341
Criconemoides lobatum: 109
Crinkle leaf: of sweet cherry, irradiation to crinkle plants produced some normal foliage, 268
Crinkle-scurf: of Valencia orange trees, genetic abnormality not transmissible but is propagatable, 31
Cristulariella pyramidalis: 747
Cronartium coleosporioides (Peridermium harknessii): 328; 455; 650
--- comandrae: 305
--- comptoniae: 538
--- fusiforme: 419; 597
--- quercuum: 328; 538
--- ribicola: 391
Crook root: of watercress, caused by Spongospora subterranea f. sp. nasturtii, 859
Cropping: 5 months' greenhouse tests on plant crops for buildup of Longidorus africanus showed greatest increase with sugarbeet, 748
Crop rotation: oat mosaic virus limiting factor in oat production when susceptible varieties are grown without rotation, 430
Crotalaria juncea: leaf blight disease of, caused by Bipolaris tetramera in India, 377
--- spectabilis: as cover crop, 573
Crown gall: Agrobacterium tumefaciens, soil fumigants tested for control of, in Oregon, 102; of raspberry, Agrobacterium tumefaciens enters only wounded plant tissue, no relationship found between pathogen and other pathogenic organisms, 492; 20 cases shown of root-knot nematode infesting crown gall tissue in Hawaii, 550
Crown rot (see Rot)
Crown rust (see Rust)
Crownvetch (Coronilla varia): Sclerotinia trifoliorum new disease of, 882
Crucifers: Brassica geniculata apparently new host for Macrophomina phaseoli in California, 223; Septoria leaf spots on wild crucifers in India, 175
---: cabbage, diseases of in Colorado in 1967, 542; stubby root nematode reached much higher populations on cabbage after soil fumigation with DD and EDB than with DBCP, 573
---: turnip, turnip mosaic virus identified in Florida, 759
Crumbly berry: of red raspberry, tomato ringspot virus cause of, 386
Cucumber (see Cucurbits)

Cucurbits: cantaloup, heat pasteurization and chemical fungicides for control of Fusarium rot, 262
---: cucumber: damping-off reduced by band treatment and surface incorporation of fungicides alone or combined with petroleum mulch, 806; diseases of in Colorado in 1967, 542; new symptom of gummy stem blight on cucumber fruit, 49
---: cucumber mosaic virus, 11; 171
---: powdery mildew: control on squash and cucumber plants with Fungicide 1991, 630; control with systemic fungicide, 754
---: squash mosaic virus and cucumber mosaic virus are most prevalent cucurbit viruses found in Kansas survey, 11; in Venezuela, 171
---: watermelon: bacterial rind necrosis, apparently first published report of occurrence on American continent, 375; downy mildew preventive spray schedule resulted in slower disease development on plants, 979
---: watermelon mosaic virus 1 and 2, cucumber mosaic virus and squash mosaic virus found in cucurbit-growing areas of Venezuela, 171
Cultures: inhibition of growth of Armillaria mellea on potato-dextrose agar slants with several systemic chemicals, 639; method of conserving cultures in viable state using Saran Wrap and polyethylene as tube closures, 961; 12 cultures of Erysiphe graminis hordei from four countries used to study interaction of 60 varieties of barley to cultures diverse in pathogenicity, 463; stem tip, to produce geranium plants, 600
Curly top: of tomatoes, local spread is restricted during low temperature periods in Washington, 17
Curvularia disease: of Gladiolus in India, 662
Cylindrocarpon tenue: 773
Cylindrocladium floridanum: 615
--- scoparium: 615
Cypress (see Baldcypress)
Cytophoma pruinosa: 105
Cytospora cincta: 154

Damping-off: of cucumber, as affected by petroleum mulch, fungicides and methods of application, 806
Datura: injury due to air pollution in Connecticut, 732
Daylily (Hemerocallis spp.): new leaf-streak disease of daylily found in Mississippi, 751
Decline: of Pinus ponderosa in southern California, apparently from prolonged exposure to aerial oxidants, 707; of shelterbelt trees in South Dakota, dagger nematode may be major factor in premature decline, 795
Deightoniella torulosa: 405
Delaware: 95; 683; 685
Delonix regia: ornamental tree of India susceptible to Pythium root rot, 705
Diaporthe vaccinii: 341
Dieback: of cranberry, isolation of Diaporthe vaccinii from diseased stems in Wisconsin, 341; of pecan caused by Botryosphaeria berengeriana in Georgia and Florida, 947; of redbud tree, caused by exhaust fumes from automobile, 744; of white ash in New York, rate of symptom development and rate of spread within given areas, 105
Diplodia gossypina: 582
--- maydis: 611
--- pinea: 359
--- tumefaciens: 829
Diplogaster lheritieri: 3
Disease control: commercial implementation in Florida effective, 339; use of surface weather maps, upper air soundings and prognostic charts should improve weather forecasts for fusiform rust control programs, 419
Diseases: of canning crops in Colorado, 542; of Florida flue-cured tobacco in 1968, 984; reduction in yield of cotton caused by, 314; of tobacco in Kentucky, 1967, 589
Distribution: distribution of cucurbit viruses in Kansas, 11; of Dutch elm disease west of Mississippi River in 1967, 652; of nematodes associated with soybeans, survey of Alabama, Florida and Georgia, 40; distribution by State of physiologic races of Puccinia f. sp. tritici and Puccinia graminis f. sp. avenae in the United States in 1965, 1966, 1967, 512; 642; 897
DMSO (see under Chemicals)
Douglas-fir (see Pseudotsuga)
Downy mildew: of corn, further evidence of the seedborne nature of two corn downy mildews and their possible control with chemicals, 446
--- ---: of grapevines, Plasmopara viticola, 372
--- ---: of lentil (Lens esculenta) in India, 817
--- ---: of lima bean, fungus has been kept in viable and pathogenic condition for 476 days at -60°C, 507; strains A and B of Phytophthora phaseoli not

pathogenic to two new lima bean lines G 1 and G 2, 563
--- ---: of sugarbeet, elimination as a major sugarbeet disease in California since more resistant varieties are grown, 297
--- ---: of watermelon, disease development slower on plants sprayed with tetrachloroisophthalonitrile prior to disease symptom appearance, 979
--- ---: of wheat in India, 569
Dutch elm disease: first known incidence of disease in Idaho, 46; recorded distribution west of the Mississippi River as of 1967, 652; South Dakota, first known occurrence, 1967, 349
Dwarf mistletoe (see Mistletoe)

Early blight (see Blight)
Eastern gall rust (see Rust)
Ecology: of Cephalosporium stripe disease of winter wheat in Washington, 590
Eggplant (see also Solanum melongena): Heterodera tabacum invasion suppressed by cellulosic soil amendments, 441
Elm: American, Fomes annosus found pathogenic to, in Ontario, 692; Cephalosporium wilt of, believed to be first report of the fungus wilt in the Midsouth, 170
---: Dutch: Ceratocystis ulmi isolated from two American elms in Idaho, first known incidence in State, 46; found in South Dakota for first time, 1967, 349; recorded distribution of disease west of Mississippi River in 1967, 652
Elsinoë piri: 277
Enation leaf curl: of tomato, graft transmissible and considered to be strain of tobacco leaf curl virus (Nicotiana 10), in India, 595
England: 799
Enzyme: pectolytic activity in Erwinia carotovora, rapid method for preparation of pectate agar medium, 167
Epidemics: Rhizoctonia solani root rot epidemic of soybeans in central Iowa in 1967, 613
Epiphytotic: of corn smut in Indiana and Illinois, heavy loss related to cool weather, 922
Eradication: citrus canker, first entirely successful eradication campaign conducted on large scale in the United States, 667
Erwinia sp.: 375
--- amylovora: 259; 698
--- aroideae: 14
--- carotovora: 167
Erysiphe spp.: 754
--- graminis f. sp. hordei: 463
Eutypa armeniacae: 751
Exhaust fumes: from automobile carport caused severe damage to redbud tree, 744

Fairy ring: of cranberry, good control with ferbam applied immediately after harvest, 87
Feeder root necrosis: of pecan, caused by Pythium and nematodes, chemicals tested for control, 577
Fern: leatherleaf (Polystichum adiantiforme): control of Pratylenchus penetrans on roots by use of nematocides, 383
Fertilizer: nitrogenous fertilizers used to hasten decomposition of fall leaf drop of apple to reduce scab infection in spring, 728; organic nitrogen reduced number of nematodes on Tifgreen Bermuda grass more than inorganic nitrogen, 46
Fir (see Abies and Pseudotsuga)
Fire blight (see Blight)
Flag smut (see Smut)
Florida: 31; 118; 339; 455; 646; 759; 844; 859; 887; 984
Fluorescent spots: in cotton fiber, associated with Aspergillus flavus, caused by conversion of kojic acid to greenish-yellow substance under influence of peroxidase in fiber, 671
Fomes annosus: 370; first report of sand pine as host of Fomes annosus, 635; incidence in Maritime Provinces of Canada, 638; two new hardwood hosts, Fraxinus americana and Ulmus americana, 692
Fraxinus americana (see also Ash): new host for Fomes annosus in Ontario, 692
Fungi: control of many genera with Fungicide 1991, 95; decay fungi not previously considered important in living trees found on young grand fir trees in northern Idaho, 489; inhibitory effects of juices of various fungi on tobacco mosaic virus, 528; kind of fungi associated with cotton boll rot and their frequency, 582; mycoflora of surface-sterilized peanut kernels in Israel, 608; spoilage of cultivated mushrooms in markets associated with, 712; use of Saran Wrap and polyethylene as tube closures for conserving fungal cultures for prolonged periods in viable state, 961
---: Armillaria mellea, control of with

systemic chemicals, 639; nematode-trapping, associated with citrus nematode, effect of nematocides upon, 851
Fungicides (see also Chemicals): band treatments with NIA 5961 or Dexon reduced damping-off of cucumber, 806; coordination product of zinc ion and maneb for control of Botrytis leaf blight, air application as effective as ground application in New York, 469; fungicide blends of 1:1 Difolatan + DCNA and 2:3 captan + Ceresan superior to single fungicides in reducing seedborne fungi of peanut seed, 240; sprayer for fungicide applications to experimental turfgrass plots, 678
---: 5-aceto-8-hydroxyquinolinesulfate (G 20072) reduced incidence of decay of Valencia oranges inoculated with Penicillium digitatum better than other compounds tested, 777
---: 2-acetyl-3-hydroxyfuran (GS-7174) proved very effective in control of decay of Barlinka grapes shipped from South Africa, 479
---: Bordeaux mixture: for control of Diplodia tip blight of pines, 359; sprays of Bordeaux mixture (8-8-100) and 7.5% phenylmercury triethanolammonium lactate (PAS) controlled Cercospora blight of junipers, 361
---: chloroneb and two 1,4-oxathiin derivatives tested for chemotherapeutic activity against Rhizoctonia solani in cotton seedlings, 399
---: 1-chloro-2-nitropropane (NIA 5961) for control of pea root rot by soil incorporation and in-furrow treatments, 72; (NIA 5961) + PCNB gave effective control of head smut of corn in Idaho, 726; (NIA 5961) and sodium p-(dimethylamino) benzene-diazosulfonate (Dexon) reduced damping-off of cucumber, 806
---: copper compounds as foliar sprays for reducing spread of halo blight of snap bean, 225
---: cuprous oxide (Yellow cuprocide) for control of Myrothecium stem necrosis of coffee seedlings, 115
---: cycloheximide thiosemicarbazone (CTS), 8-quinolinol benzoate or dimethyl sulfoxide (DMSO) undiluted, prevented Cytospora infection in 4-year-old 'Italian' prune trees, 154; cycloheximide and DCMOD best of several tested for reduction of number of galls producing telia of cedar-apple rust on juniper, 692; cycloheximide and phytoactin ineffective as chemical controls of blister rust disease, 6
---: DCNA superior to mercurous chloride or PCNB soil treatments for control of onion white rot in Washington, 272
---: DMTT and SMDC applied as drench most effective of 10 chemicals in penetrating soil and killing hyphae of Phymatotrichum omnivorum in soil cultures, 232
---: dodine (Cyprex-85W) and Fungicide 1991 gave good control of pecan scab in Georgia, 108
---: ferbam, applied immediately after harvest gave good control of fairy ring disease of cranberry in Massachusetts, 87; ferbam gives effective control of fusiform rust in pine nurseries, more frequent sprayings necessary when weather conditions are favorable, 597
---: Fungicide 1991 (1-(butylcarbamoyl)-2-benzimidazole carbamic acid methyl ester): new fungicide shows preventive, curative, and systemic fungicidal properties as well as a mite ovicide effect, 95; has shown chemotherapeutic activity against several widely different grass diseases in greenhouse tests in Oregon, 205; gave outstanding control of Sclerotinia dollar spot on 29 varieties of bentgrass turf in Pennsylvania, 410; reduced root rot of pea in Alberta when applied as in-furrow treatment, 565; only one of 16 tested that gave 100% control of Cylindrocladium blights on azalea, 615; provided systemic control of powdery mildew on squash and cucumber, 630; effective in control of stripe smut in 'Merion' Kentucky bluegrass turf in New Jersey, 635; completely inhibited perithecial development of apple scab after single fall application, superior to other fungicides tested, 654; outstanding as preharvest spray for control of peach brown rot in Georgia, 695; gave long-lasting field control of brown rot on stone fruits in California, 722; as root application gave excellent control of several powdery mildews, 754; effective against Rhizoctonia solani on cotton seedlings, 781; best of three for control of postharvest decay of citrus applied as preharvest spray, 844
---: Karathane 1/4 lb, zinc + maneb 1-2 lb, plus 12 oz Triton B-1956, to 100 gal. water per acre controlled powdery mildew of papaya, 816
---: methyl bromide: effect of fumigation on mycorrhizae and growth of sweet-

gum seedlings in Mississippi, 483; methyl bromide and chloropicrin (MBR-CP) reduced soilborne wheat mosaic virus incidence in 'Pawnee' wheat and increased yield in Kansas, 559
---: nickel-maneb mixture superior to zineb or maneb in control of oat crown rust in Iowa, 205
---: 1,4-oxathiin derivative (DCMOD) applied in-furrow at planting appears to increase effectiveness of late-season sprays for rust control on wheat, 856
---: sodium pentachlorophenate (SPCP) and Bordeaux mixture as post-leaf-fall and pre-leaf-fall protective fungicides against European canker of apple, 227
---: tetrachloroisophthalonitrile: reduced Aspergillus flavus in soil in peanut study, had no effect on dominant genera of soil fungi, 718; for control of downy mildew in watermelon, 979
---: thiabendazole: thiabendazole and Fungicide 1991 more effective against Verticillium in cotton with surfactant Tween 20, 944; thiabendazole, maneb + zinc and captan effective in control of stem-end rot of cantaloup for postharvest treatment, 262
---: Trizone one of several soil fumigants tested for control of crown gall of mazzard cherry, 102
---: several tested together: disulfoton, zinc + maneb, atrazine and propazine associated with increased grain yield in corn plots inoculated with maize dwarf mosaic virus, 545; for control of anthracnose and early blight of tomato, 685; to control cotton boll rot, simple evaluating technique, 835
Furcaspora pinicola: 486
Fusarium spp.: 213; 565; 582
--- graminearum: 79; reported for the first time in the United Arab Republic, as causing seed rot and seedling blight of corn, 84
--- moniliforme: 79; 84
--- oxysporum: 939; 961
--- --- f. cubense: 901
--- --- f. sp. lycopersici: 553
--- --- perniciosum: 79; 949
--- --- f. pisi race 2: 72
--- --- f. vasinfectum: 910
--- roseum: 262; 840
--- solani: 262; 628; 939
--- wilt (see Wilt)
Fusicladium effusum: 108
Fusicoccum sp.: 105

Gall rust (see Rust)
Georgia: 108; 114; 179; 320; 334; 467; 635; 747; 947
Geranium: successful use of apical meristem and callus cultures to produce geranium plants, 600
Germination: anaerobic treatment of barley seed to kill loose smut fungus, reduced germinability during first year after treatment, 792
Gladiolus: Curvularia disease of in India, 662
Gloeocephalus hemerocalli: 751
Godronia fuliginosa: 167
Grand fir (see Abies)
Grape: 2-acetyl-3-hydroxyfuran (GS-7174) a vapor-phase chemical very effective in controlling decay of Barlinka grapes during storage and transit, 479; dagger nematodes, Xiphinema index, found infecting grapevines in Iraq, 626; limb dieback or gummosis in apricot caused by Eutypa armeniacae, found in grape vineyards in California, 751; survey of grapevine diseases in northeast Mexico, 372
Grasses: occurrence in Cochliobolus carbonum of capacities to blight gramineous hosts, 605; 150 isolates of Cochliobolus heterostrophus tested for capacity to induce blight on 34 gramineous test species, 879; several different grass diseases controlled by systemic Fungicide 1991 in greenhouse tests in Oregon, 205
---: Agropyron repens, Agropyron mosaic virus present in Montana, based on serological, host range and virus morphological evidence, 139
---: bentgrass, varietal susceptibility to Sclerotinia dollar spot and control with Fungicide 1991, 410
---: Bermuda grass: activated sewage sludge (organic nitrogen) more effective than inorganic nitrogen in reduction of sting nematodes in turf of 'Tifgreen' Bermuda grass, 46; nematode Hypsoperine graminis found in Virginia on 'Tifgreen' Bermuda grass, 162
---: bluegrass: 'Merion' Kentucky bluegrass, relative amounts of stripe and flag smut incidence in Ohio, 633; stripe smut on 'Merion' Kentucky bluegrass effectively controlled with Fungicide 1991 in New Jersey, 635
---: broomcorn, and Sudan grass, maize dwarf mosaic virus found on, in New Mexico for the first time, 365
---: turfgrass: damage to several turfgrasses from ring nematode, Criconemoides lobatum, caused mainly by reduction of absorptive surface of roots, 109; plot sprayer (carbon

dioxide-powered) for turfgrass disease control fungicide application, 678
Gray mold: of grapes, control of in Barlinka grapes during storage and transit, 479; of green onions caused by Botrytis allii, 974
Green mold: of Valencia oranges, in screening trials 5-aceto-8-hydroxyquinolinesulfate showed some promise for control, 777
Guatemala: 115
Gummosis: or limb dieback of apricot found on grape, Eutypa armeniacae causal fungus, 751
---: of citrus, caused by Phytophthora spp., some factors in resistance, 952
Gymnosporangium juniperi-virginianae: 692

Hardwoods: nematode associations since 1961 with southern hardwoods and conifers, 837
Hawaii: 56; 405; 515; 550; 814; 946
Head smut (see Smut)
Helicotylenchus dihystera: 150
Helminthosporium maydis: 134
--- pedicellatum: 477
--- rostratum: 784
--- turcicum: 427
Hemlock (see Tsuga)
Hericium abietis: 489
Heterodera sp.: 626
--- spp.: key to the Heterodera "groups", 827
--- cacti: 114
--- glycines: 45
--- punctata: 479
--- rostochiensis: first infestation of golden nematode on mainland of North America, in western New York, 184; 587
--- sacchari: 237
--- tabacum: 57; 441
Hevea brasiliensis (rubber tree): recovery of Phytophthora spp. from infected Hevea bark with unripe cacao pod, 5
Hibiscus cannabinus (kenaf): several common plant viruses pathogenic to fiber plant, 530
Hibiscus esculentus (okra): Alternaria tenuis isolated from leafspots on new plant introduction, 248; reduction of root-knot disease obtained with use of organic amendments over 4-year period in outdoor microplots in California, 315
--- rosa-sinensis: foliar nematode found on, in Florida, 118
Honduras: 901
Hoplolaimus columbus: 571
Hops: Phytophthora crown and root rot, wilt phase in several yards in Sacramento County, California, 76
Horse: for production of large quantities of antiserum of tobacco mosaic virus, 860
Humidity: modified microhumidity chamber for pathogenicity studies, 327
Hyalopsora aspidiotus: 336
Hydrogen-ion: influence of concentration on germination of oospores of Aphanomyces euteiches, 264
Hymenochaete agglutinans: 496
Hypoxylon pruinatum: 659
Hypsoperine graminis: 46; 162

Idaho: 46; 295; 965
Illinois: 68; 223; 692
India: 145; 175; 300; 377; 436; 477; 494; 626; 773; 784
Indiana: 223; 395
Indonesia: 122
Infectivity: of Meloidogyne arenaria after ingestion by bovine in Rhodesia, 99
Inhibition: inhibitory effects of juices of various fungi on tobacco mosaic virus, 528
Inoculum: bacterial canker of tomato, level necessary for infection, 741
---: concentration, of sugarcane mosaic virus, screening seedlings for levels of resistance to strain H of virus, 705
In-vitro: geranium plants differentiated in-vitro from stem tip and callus cultures, 600
Iowa: 59; 205; 459; 613
Iran: 687
Iraq: 626
Isatis tinctoria: seed oil plant introduction heavily infected with Colletotrichum higginsianum, 248
Israel: 310; 472; 525; 608; 718

Juniperus spp.: Cercospora blight, occurrence and control in Nebraska, 361; plant-parasitic nematodes associated with in New Mexico, 52
--- scopulorum: control studies on Gymnosporangium juniperi-virginianae, 692
--- virginianae: fungicides tested for control of cedar-apple rust, 692

Kansas: 11; 357
Kentucky: 71; 204; 589
Key: to Heterodera "groups" to separate and identify the golden nematode, Heterodera rostochiensis, 827

Leaf: pathogenicity studies with use of microhumidity chamber, 327
Leaf blight (see Blight)
Leaf scald: of sugarcane in Florida, first

report in continental United States, 646
Leaf spot: several plant introductions in SIPE program found susceptible to common leafspot fungi, 248; of celery, caused by Septoria apiicola, 142; of clusterbeans (Cyamopsis tetragonoloba), occurrence of new Alternaria leaf spot of clusterbeans in India, 145; of peanut, 10 varieties tested for susceptibility to Cercospora personata in Sudan, 405; of pecan, zonate-leafspot caused by Cristulariella pyramidalis, 747; on tomato and potato in Hawaii, caused by Phoma destructiva, first confirmation of foliar infection, 946; two new Septoria leaf spot diseases on Indian wild crucifers, 175
Leaf-streak: of daylily, caused by previously undescribed fungus, Gloeocephalus hemerocalli, 751
Legumes: diseases of pulse crops in India in 1966, 300; diseases of pulse crops (edible legumes) in Iran, 687
Lentil (Lens esculenta): downy mildew, apparently first record of pathogenicity to lentils, 817
Lentinellus montanus: 820
Limb dieback: of apricot found on grape, 751
Limnanthes douglasii: potential seed oil plant infected by several common plant viruses, 530
Little leaf disease: of Solanum melongena in India, severity of disease calculated by two methods, one based on the number of diseased and healthy leaves of diseased plants, and the other based on the number of healthy leaves of diseased and healthy plants, 118
Longevity (see also Survival): Heterodera glycines larvae in cysts when stored in seed bags remained infective 17, 18, and 22 months, 45; Meloidogyne arenaria eggs or larvae still infective after ingestion by a bovine, 99; Puccinia recondita, effect of temperature on viability of urediospores on dry wheat foliage, 186; survival of Cercospora kikuchii on soybean stems in the field, 931; survival time of exocortis virus of citrus on contaminated knife blades, 935; test tube slants of Fusarium oxysporum stored at 5°C with plastic closures viable after 35 months' storage, 961
Longidorus africanus: 748
--- vineacola: 525
Losses: Fusarium stem rot of carnation, in commercial production of cuttings, 840
Louisiana: 120; 189; 307; 449; 667; 775; 780
Luffa acutangula: new host for tobacco ringspot virus, 374

Macrophomina phaseoli: 223
Malaysia: 5; 132
Malva parviflora: apparently a new host for Macrophomina phaseoli in California, 223
Manitoba: 829
Maple: big leaf maple and silver maple hosts for European mistletoe, Viscum album, in California, 198
Market diseases: of corn, southern corn leaf blight on sweet corn, 847; of mushrooms, microorganisms associated with market deterioration, 712; of onion, gray mold of green onions, 974
Maryland: 259; 284
Massachusetts: 87
Medium: modified Goto's medium to induce sporulation of Piricularia oryzae, rice blast pathogen, 703
Meloidogyne spp.: 550
--- arenaria: 99
--- graminicola: 423
--- hapla: 492; 956
--- incognita: 315
--- javanica: 150
Methods and techniques: cotton boll rot, method for preliminary evaluation of large numbers of systemic compounds for control, 835; estimating severity of little leaf disease of Solanum melongena, 118; formulae presented for calculating crop loss in yield in a locality or region, 963; modification of cyclone collector for collecting rust fungus urediospores directly into vials with addition of spore guide tube, 148; modified Petri dish microhumidity chamber, 327; sampling of air spora to detect Fomes annosus incidence in Maritime Provinces by Rishbeth method, 638; selective staining of dead cells of ozone-injured bean leaves, method useful to detect leaf cells killed by any agent or pathogen that does not digest starch grains of injured cell, 732; specially designed chamber to test effect of temperature on response of plants to ozone, 957; supporting plants in nutrient solution, 158
---: antiserum: rapid serodiagnostic test for potato ring rot, 502; immunization of horse to produce tobacco mosaic virus antiserum in large quantity, 860
---: cultures: cacao pod used for recovering Phytophthora species pathogenic

to Hevea brasiliensis, 5; preparation of a semi-solid agar medium for detection of pectolytic enzyme activity in Erwinia carotovora, 167; isolation of Sclerotium cepivorum in pure cultures by effective and simple method, 277; glass dish, flask, and clay pot techniques to determine chemotherapeutic activity of certain fungicides against Rhizoctonia solani in cotton seedlings, 399; system for rapid classification of Puccinia recondita tritici cultures, 437; plastic film closures in the conservation of fungal cultures, 961
---: inoculation: air-blast inoculation, plot techniques with maize dwarf mosaic virus: I. field studies, 1966, 24; air-blast inoculation, plot techniques with maize dwarf mosaic virus: II. greenhouse studies, 1966, 28; artificial inoculation by toothpick method at base of plant superior to foliar applications of conidia, in differentiating for susceptibility to anthracnose among sorghum lines, 164; mechanical planter for system of rapid classification of Puccinia recondita tritici cultures and a method of data processing suitable for computer analysis, 437; CO_2-powered BB pistol promising technique for artificial inoculation of corn plants for reaction to Diplodia stalk rot, 611; stem and root inoculation methods for testing cacao clones for resistance to Phytophthora palmivora, 790
---: spray or fumigation: apparatus for subsurface application of nematocides to nursery plants in containers, 218; garden tractor applicator for fumigation of small plots, 575; specially designed sprayer for fungicidal applications in experimental turfgrass plots, 678
Mexico: 372
Michigan: 37
Microscopy: electron, of newly described virus, Bidens mottle virus, 763
Microxyphiella hibiscifolia: 602
Millet (see Sorghum)
Mimosa: Fusarium wilt can be seedborne, 949; vascular disease caused by Fusarium oxysporum perniciosum found in Arkansas, 79
Minnesota: 328
Mississippi: 170; 483; 582; 751; 822
Missouri: 739; 811
Mistletoe: dwarf: lodgepole pine dwarf mistletoe on ponderosa pine in Colorado, 125; infection of Scots pine in Canada by Arceuthobium americanum, 409; Arceuthobium campylopodum infection on white fir, 452
---: eastern, Phoradendron flavescens found in Alabama on water oak being girdled by Hymenochaete agglutinans, new record of Phoradendron as host for pathogen, 496
---: European, Viscum album rediscovered in California on apple and several other host trees, 198
Monilinia fructicola: 722
Monilochaetes infuscans: 475
Montana: 139; 184; 201; 818
Musa sp.: disease resistant accession 'Pelipita' from Philippines found as substitute for fusarial and bacterial wilt susceptible 'Bluggoe' and ABB Musa, 901
Mushroom: microorganisms associated with market deterioration of cultivated mushroom, Agaricus bisporus, 712
Mycosphaerella melonis: 49
Myrothecium roridum: stem necrosis and leaf spot, an uncommon disease of coffee in Guatemala, 115

Nebraska: 357
Necrosis: cotyledon, in soybean, 459
Nectarine (see Prunus)
Nectria galligena: 227
Needle cast: caused by Bifusella linearis found on white pine, 305
--- ---: caused by Rhizosphaera kalkhoffii on blue spruce, 804
Needle rust (see Rust)
Nematocides: a combination of specific pesticides as opposed to broad-spectrum soil fumigants for multiple pest control in Georgia, 193; applicator mounted on a tractor designed for fumigation of small plots, 575; preplanting soil fumigation superior to postplanting fumigation in California peach, walnut and prune orchards, 890
---: cynem, subsurface drench with cynem and similar compounds more efficient than surface drench for nursery plants in containers, 218
---: DBCP: superior to DD or EDB in residual action following soil fumigation to control Trichodorus christiei in Florida, 573; DBCP fungistatic but not lethal to nematode-trapping fungi associated with citrus nematode, several nematocides tested, 851
---: ethylene dibromide (EDB), as single preplanting treatment gave excellent control of nematodes on onion in loessial soils in Israel, 525
---: several tested together: soil treatment with O,O-diethyl S-(ethylthio)=

methyl phosphorodithioate, O,O-diethyl O-((p-methylsulfinyl)phenyl) phosphorothioate, and 2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl)oxime controlled sting and stubby-root nematodes and promoted good growth of vegetable crops in Georgia, 19; in-the-row treatment with some nematocides in the halogenated hydrocarbon group and some new nematocides belonging to organic phosphate and carbamate groups gave good control of reniform nematodes on sweetpotatoes in Louisiana, 127; cynem + phorate, 2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl)oxime (UC 21149) and 2,3-dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate (N-10242) controlled Pratylenchus penetrans on leatherleaf fern, 383; four dichloropropene fumigants significantly reduced the soil population of Pratylenchus penetrans in soil of high organic content in Indiana, 519; DBCP, 1,3-dichloropropene, 2-methyl-2-(methylthio) propionaldehyde O-(methylcarbamoyl)oxime (UC 21149) and 1,3-dichloropropene and 1,2-dichloropropane mixture (DD) controlled reniform nematodes on cotton, 786

Nematodes: burrowing, Radopholus similis associated with decline in young tea fields in Ceylon, first record of nematode in Ceylon, 528

---: citrus, several nematocides tested for effect on nematode-trapping fungi associated with Tylenchulus semipenetrans, 851

---: Columbia lance nematode, Hoplolaimus columbus, associated with cotton and soybeans in South Carolina, 571

---: cyst (see also golden): key to Heterodera "groups", 827; of cactus, first record of Heterodera cacti in Georgia, 114; of grass, first report of occurrence of Heterodera punctata in Rhode Island, 479; Heterodera sp. found in association with sugarcane for the first time in India, 626; soybean cyst nematode, larvae in cysts remained infective when stored in seed bags up to 22 months and longer, 45; of sugarcane, Heterodera sacchari first collected in Nigeria in 1965, 237; tobacco cyst nematode, cellulose soil amendments and fertilizers suppressed nematode invasion of eggplant, 441

---: dagger: nematodes on grapevines in Iraq, first record, 626; nematode association with premature decline of shelterbelt trees in South Dakota, 795

---: foliar: disease of tuberose caused by Aphelenchoides besseyi, 56; Aphelenchoides fragariae found infesting Hibiscus rosa-sinensis in Florida, 118; on aquatic plants, 455

---: golden nematode: in potato field in western New York, first report on mainland of North America, 184; discovered in potato-growing region of Cerro Punta, Panama, 587; Heterodera "groups" and species within Heterodera rostochiensis "group" keyed to simplify indentification of golden nematode, 827

---: needle: Longidorus africanus, effect of cropping on population levels, 748; Longidorus vineacola controlled on onion in Israel by single preplanting treatment with ethylene dibromide, 525

---: plant-parasitic: in central New Mexico associated with marginal Pinus ponderosa stands, 52; susceptibility of parasitic nematodes to sub-freezing temperatures, 768; geographic tabulation of new plant-parasitic nematode associations with southern forest hardwoods and conifers other than pines, 837

---: pseudo-root-knot, Hypsoperine graminis found in 'Tifgreen' Bermuda grass in Virginia, 162

---: reniform: several nematocides tested for control of Rotylenchulus reniformis, 786; on sweetpotato, evaluation of several nematocides during 5-year period in Louisiana, 127

---: ring, on turfgrasses, principal damage caused by reduction of fibrous roots, 109

---: root-knot: simultaneous occurrence of root-knot nematodes and crown gall bacteria, 550; Meloidogyne arenaria infective after passage through digestive tract of cattle fed Meloidogyne-infected potato tubers in Rhodesia, 99; Meloidogyne graminicola, a new pest of rice, common name of rice root-knot nematode proposed, 423; Meloidogyne hapla showed difference in degree of galling of potato varieties but all varieties susceptible, 956; Meloidogyne incognita, infectivity and survival reduced by organic soil amendments, 315; Meloidogyne naasi discovered in Oregon on barley, 169

---: root lesions: Pratylenchus penetrans pathogenic to leatherleaf fern, effective control with nematocides, 383; reduction of soil population of Pratylenchus penetrans by use of four di-

chloropropene fumigants in Indiana, 519
---: saprozoic nematodes transmit Streptomyces phage, 3
---: sting, Belonolaimus longicaudatus, found pathogenic to sycamore tree seedlings in Georgia, 523
---: sting and pseudo-root-knot, on 'Tifgreen' Bermuda grass, organic nitrogen fertilizer reduced number of nematodes more than inorganic nitrogen, 46
---: sting and stubby-root, organophosphate and carbamate pesticides gave effective control in Georgia, 19
---: stubby-root: re-establishment of Trichodorus christiei after soil fumigation in Florida, 573
---: several together: associated with soybeans in Alabama, Florida and Georgia, 40; mutual antagonism between Heterodera tabacum and some other parasitic nematodes, 57; Xiphinema spp., Tylenchorhynchus spp., and Trichodorus spp., nematodes most frequently collected around declining rose plantings in United Arab Republic, 149; Helicotylenchus dihystera, Meloidogyne javanica, and Xiphinema elongatum associated with olive decline in United Arab Republic, 150; effort to relate high population of nematodes to high Aphanomyces root rot disease from Wisconsin pea fields unsuccessful, 190; root-knot, spiral and burrowing, control in plant containers with use of applicator for subsurface drench, 218; associated with pecan tree decline in Georgia, 334; associated with grapevines in Mexico, 372; Telotylenchus loofi, Aphelenchus avenae, and Ditylenchus myceliophagus most widely distributed in soil samples from 80 fields of great millet in India, 396; 104 host-nematode associations are new records for Hawaii, 53 thought to be new records for the host-nematode association, 515; stubby-root, root-knot and lesion nematodes associated with feeder root necrosis of pecan, 577; possible role in transmission of cherry raspleaf virus, 618; reduction with preplant and postplant soil fumigation in peach, walnut and prune orchards, 890
Net blotch: of barley seedlings, excellent resistance to Pyrenophora teres substantiated, 215
New diseases: Bidens mottle virus isolated and identified in Florida from Bidens pilosa and Lepidium virginicum, 763; leaf-streak of daylily, fungus not previously described, 751; on Phaseolus mungo, P. aureus and Cajanus cajan in India, 300; Sclerotinia trifoliorum caused chlorosis, wilting and death of crowns in crownvetch, 882; stem pitting of peach, 91; 287; 292; 907; vascular wilt caused by Verticillium nigrescens found in Australia, 904
New distribution: list of fungus pathogens not previously reported from Chile, 703
---: alfalfa anthracnose in Washington, 983
---: alfalfa mosaic virus on pepper and tomato bushy stunt virus on tomato for the first time in Argentina, 674
---: bacterial diseases: bacterial canker of tomato in Argentina, 595; bacterial rind necrosis of watermelon in Texas, 375; fire blight distribution in the world, confirmed in Poland, Netherlands, and Guatemala in 1966, 698
---: boll rot of cotton caused by Phytophthora sp. in Louisiana, 780
---: brown spot of tobacco, first occurrence in Kentucky in certain dark and burley tobacco-growing areas, 1966, 71
---: Cephalosporium wilt of American elm, believed to be first report of in the Midsouth, 170
---: crook root of watercress in Florida and Pennsylvania, has previously been reported only from Europe, 859
---: Deightoniella torulosa on banana in Hawaii, first published record of occurrence, 405
---: Dutch elm disease: first known occurrence in South Dakota, 1967, 349; in Idaho, 46
---: Eutypa armeniacae found infecting grape wood, 751
---: exocortis virus in Ortanique orange in Jamaica, 741
---: Fusarium graminearum and Nigrospora oryzae cause seed rot and seedling blight of corn in United Arab Republic, first report, 84
---: Fusarium oxysporum f. sp. lycopersici race 2 in New Jersey tomato-growing region, 553
---: Helminthosporium pedicellatum on corn in India, 477
---: leaf scald of sugarcane found in Florida in 1967, 646
---: legume diseases in Iran, many thought to be previously unreported, 687
---: maize dwarf mosaic virus identified in New Mexico for the first time in

1967, 365
---: needle disease, Furcaspora pinicola, found in Humboldt Co., California infecting dead needles of Abies concolor, 486
---: nematodes: new records for 104 new host-nematode associations for Hawaii, 515; new nematode associations with southern hardwoods and conifers other than pines since 1961, 837
---: ---: cyst: cactus cyst nematode in Georgia, 114; cyst nematode, Heterodera sacchari, first collected around sugarcane roots in Nigeria, 1965, 237; grass cyst nematode, Heterodera punctata, in Rhode Island, 479
---: ---: golden: found in western New York, 184; found in Panama, first report of, 587
---: oak wilt in Platte County, Missouri, 811
---: sooty mold of cotton, caused by Microxyphiella hibiscifolia, in India, 602
---: Verticicladiella wagenerii found for first time in El Dorado Co., California infecting Douglas-fir, 393
---: white pine blister rust found on foxtail pine in California, 391
---: white pine needle cast caused by Bifusella linearis in California, 305
---: zonate-leafspot of pecan in Alabama and Georgia, 747
New hosts: Alternaria leaf spot caused by Alternaria temuissima on clusterbeans, new record on host, in India, 145
---: Armillaria mellea on carnation in California, 415
---: Colletotrichum sp. on pineapple in India, 436
---: crown gall, Ardisia crispa, Begonia coccinea, and Grevillea robusta, new hosts in Hawaii, 550
---: Cylindrocarpon tenue, root rot disease of coffee found in India, 773
---: European mistletoe, Viscum album, apparently first record on big leaf maple, 'Santa Rosa' plum and California buckeye in California, 198
---: Fomes annosus, sporophores found on Ulmus americana and Fraxinus americana in Ontario, 692
---: Furcaspora pinicola on dead needles of white fir in California, 486
---: Fusarium solani associated with stem canker and pink stain of teak, 628
---: Godronia fuliginosa on Populus tremuloides (trembling aspen) in Quebec, 167
---: Helminthosporium rostratum caused leaf blight of papaya in India, 784
---: Macrophomina phaseoli on Malva parviflora and Brassica geniculata, apparently new hosts, 223
---: needle cast fungus found on Engelmann spruce in Arizona, 804
---: nematodes: host-nematode associations, possibly 53 new records, from Hawaii, 515
---: ---: dagger nematodes on grapevines in Iraq, 626
---: ---: foliar: Aphelenchoides besseyi on tuberose in Hawaii, 56; A. fragariae on aquatic plants in Florida, 455; on Hibiscus rosa-sinensis, 118
---: ---: Pseudo-root-knot, Hypsoperine graminis, reported for first time infecting Agrostis palustris (C-7), Dactylis glomerata and Hordeum vulgare, 162
---: ---: root-knot, Meloidogyne javanica, in Ardisia crispa and M. incognita acrita in Grevillea robusta in Hawaii, 550
---: ---: spiral, Scutellonema brachyurum, as parasite of soybean, first report, 40
---: Pestalotia longi-setula on strawberry fruit, cause of fruit rot in Israel, 472
---: Pythium myriotylum pathogen of rye, wheat, oats and ryegrass, and cause of stem rot of tomato, in Georgia, 179; apparently first report of P. myriotylum attacking snapbeans under field conditions in the United States, in Georgia, 416
---: Septoria leaf spots on two wild crucifers in India, 175
---: "strangling fungus" Hymenochaete agglutinans as parasite of eastern mistletoe, 496
---: tobacco ringspot virus on Luffa acutangula, 374
---: white pine blister rust on foxtail pine for first time, 391
---: white pine needle cast on Pinus flexilis and P. albicaulis, 305
---: for Verticillium nigrescens in Australia, 904
---: zonate-leafspot disease on pecan, 747
New Jersey: 496; 553; 635
New Mexico: 52; 365
New York: 105; 184; 235; 268; 469; 865
New Zealand: 142
Nigeria: 134; 136; 237
Nigrospora oryzae: 79; reported for the first time in the United Arab Republic, as causing seed rot and seedling blight of corn, 84
North Carolina: 344; 528; 895

Nuclear behavior: and aeciospore germ tube morphology as basis for identification of Peridermium harknessii in eastern Canada, 650
Nutrients: method for supporting plants in nutrient solution, 158; susceptibility to Fusarium wilt in alfalfa greatest at low levels of N, P, K, or Ca, and decline with increased concentrations, 939

Oak: host of Phorodendron flavescens in turn parasitized by Hymenochaete agglutinans, 496; root rot fungus, systemic chemicals inhibited growth of Armillaria mellea on culture medium, 639
---: wilt: Ceratocystis fagacearum found in Platte County, Missouri, 811; distribution in Nebraska and Kansas, 357
Oatgrass: tall, resistance to crown rust found in 16 selections from 14 plant introductions in field and greenhouse tests in New York, 235
Oats: crown rust: 130 strains of oats screened for tolerance to common races of crown rust, 13 found superior to Cherokee, 59; mixture of $NiSO_4$ and maneb gave better control of oat crown rust than either zineb or maneb, 205
---: pathogenicity of several Pythium species to oat, barley and wheat seedlings, 209
---: stem rust: Puccinia graminis avenae, distribution and prevalence of physiologic races in the United States in 1965, two new races found, 512; race 6AF most common race in United States in 1966, 897; also in 1967, 642
---: virus, oat mosaic virus is apparently limiting factor in production when highly susceptible varieties such as Victorgrain 48-93 are cropped without benefit of rotation, 430
Odontia bicolor: 489
Ohio: 633
Oidium sp.: 754
--- caricae: 814
Oil: to control ripe fruit rots in tomato, 555
Oklahoma: 14; 158
Okra (see Hibiscus esculentus)
Olive: nematodes thought to be primary inciting agents of olive decline in United Arab Republic, 150
Olpidium: 379
Onion: Botrytis leaf blight, control by aircraft application with protective fungicide, 469
---: gray mold of green onions on the market caused by Botrytis allii, 974
---: nematodes: control effective in Israel with treatment of ethylene dibromide (EDB), onion yields increased 160%, 525; factors influencing population fluctuation of Pratylenchus penetrans in soils of high organic content, effect of four dichloropropene fumigants, 519
---: white rot, control by means of soil chemicals in southeastern Washington, 272; isolation of pathogen, Sclerotium cepivorum, in pure cultures, 277
---: virus, mechanical transmission of onion yellow dwarf virus to mother bulbs with contaminated knife, 581
Ontario: 659; 692
Ophiobolus graminis: 716
Oregon: 169; 205
Ozone: control of Penicillium molds on oranges and lemons in storage, 245; damage to apples from exposure to high rates, in Connecticut, 730; effect of temperature on response of plants to ozone in specially designed plant fumigation chamber, 957; selective staining of dead cells in bean leaves injured by ozone, 732; thought to be cause of decline of Pinus ponderosa in southern California, 707

Pacific Northwest: increase in incidence of take-all of wheat in irrigated fields, 716
Panagrellus redivivus: 3
Panama: 587
Papaya: a new leaf disease caused by Helminthosporium found in India, 784; powdery mildew control with fungicides, 814
Patch canker: of rubbertree, Hevea brasiliensis, caused by Phytophthora palmivora, 132
Pathogenicity: Godronia fuliginosa found pathogenic to aspen as well as willow, 167; of Phacidium taxicolum on Canadian yew, 403; of Scleroderris lagerbergii, additional information from Canadian tests, 738; of sting nematode on sycamore, 523
Pathogens: plant, additional list of pathogens in Chile, not previously reported, 703
Pea: diseases of canning peas in Colorado in 1967, 542; disease reaction of plant introductions to three legume viruses, 160
---: root rot: Aphanomyces, hydrogen-ion concentration influences germination of oospores of root rot fungus, 264; attempt to establish relationship

of nematodes to Aphanomyces root rot potential in Wisconsin unsuccessful, 190
---: ---: Fusarium, control by soil-incorporation and in-furrow treatments with 1-chloro-2-nitropropane, 72; Fungicide 1991 as an in-furrow application reduced Fusarium and Rhizoctonia root rot of garden pea and increased yield, in Alberta, 565
---: sweet, powdery mildew, systemic fungicide gave excellent control, 754
Peach (see also Prunus): brown rot, Fungicide 1991 best of several fungicides tested for control of Rhizopus rot, 695
---: nematode control studies with preplant and postplant fumigation, 890
---: stem pitting: widespread distribution in several Middle Atlantic States, cause yet unknown, 91; of peach, apricot and nectarine seedlings, 287; a serious new disorder of peach trees in Pennsylvania, 292; known to exist in only one orchard in South Carolina, 907
Peanut: seed mycoflora, germination, and seedling emergence as affected by seed treatment fungicides, 240; 10 varieties tested in Sudan for susceptibility to Cercospora personata, MH 383 and Baladi Bunch least susceptible, 405; 35 species of fungi survived surface disinfection with mercuric chloride, 0.1%, 608; effect of fungicides on development of fungi in soil and on kernels, 718
---: virus, new record of virus disease of peanut in India, name of "bud necrosis" proposed, 494
---: peanut mottle virus, differences in resistance indicated, but no immunity found in 465 introductions screened in Georgia, 467
---: stunt virus: not transmitted from infected white clover to peanuts in South Carolina, 926; seed transmission of virus through symptomless plants found very low, 522; varietal reaction of 300 varieties and breeding lines to, 914
Pear: fire blight, caused by Erwinia amylovora, its recent spread and present distribution, 698
---: virus: pear decline virus, leaf drop symptom associated with, 363; pear leaf curl virus transmitted by pear psylla, Psylla pyricola, 924
---: ornamental, Coniothyrium canker on, in Washington, D. C. area, 792
Pecan: varieties tested for resistance to scab and bunch disease in Louisiana, 307; unrecorded zonate-leafspot disease found in Alabama and Georgia caused by Cristulariella pyramidalis, 747
---: dieback, in Teche variety caused by Botryosphaeria berengeriana, 947
---: feeder root necrosis, caused by Pythium spp., and nematodes, treated with fungicides, 577; role of nematode and Pythium species to, 334
---: scab, good control with dodine-85W and Fungicide 1991, 108
---: shuck disease, adverse conditions which cause loss of effective leaf area favor shuck disease, 189
Peck: of cypress, caused by Stereum taxodii, 482
'Pelipita' (ABB type of Musa): Philippine accession found resistant to fusarial and bacterial diseases to eventually replace disease-susceptible 'Bluggoe' and ABB Musa, 901
Pellicularia sasakii: 323
Penicillium spp.: 245
--- digitatum: 623
--- expansum: 865; 867
--- italicum: 623
--- oxalicum: 79; 84
Pennsylvania: 24; 28; 292; 545; 859; 972
Peperomia: crown rot, first report of Phytophthora nicotianae parasitica as pathogen, 244
Pepper: diseases of cherry pepper in Colorado in 1967; virus resistance, 381
---: bacterial spot, several chemicals tested for control of Xanthomonas vesicatoria in Delaware, 683
--- viruses: alfalfa mosaic virus reported for the first time on, in Argentina, 674; varietal reaction to naturally occurring viruses in California, 928
Peridermium harknessii (Cronartium coleosporioides): 455; 650
Peronospora sp.: 817
--- schachtii: 297
--- tabacina: 310
Persimmon: wilt, vascular disease caused by Cephalosporium diospyri found in Arkansas, 79
Pestalotia longi-setula: 472
Petunia: air pollution damage, 732
Phacidium taxicolum: 403
Phage: Streptomyces griseus phage can be transmitted by saprozoic nematodes, 3
Pholiota adiposa: 489
Phoma destructiva: 946
Phoradendron flavescens: possibly first report that Hymenochaete agglutinans is a parasite of P. flavescens (eastern mistletoe), 496

Phymatotrichum omnivorum: 232; 372
Physiological diseases: cotyledon necrosis of soybean, 459; shuck disease of pecan, appears to be caused by a physiological-nutritional imbalance, 189
Phytophthora spp.: 5; 76; 780; 952
--- cryptogea: 280
--- megasperma sojae: 355
--- nicotianae parasitica: first report as pathogen of crown rot of Peperomia, 244
--- palmivora: 5; 132; 790
--- parasitica: 284
--- phaseoli: 507; 563
Picea spp.: pathogenicity of Scleroderris lagerbergii, continued study in Quebec, 738
--- engelmannii (Engelmann spruce): butt rot caused by Lentinellus montanus, 820; new host for needle cast fungus, Rhizosphaera kalkhoffii, 804
--- pungens (blue spruce): needle cast fungus, Rhizosphaera kalkhoffii found in native blue spruce stands, 804
Pineapple: new spot anthracnose disease of pineapple caused by Colletotrichum sp. in India, 436
Pink rot: of potatoes in Western Australia, 280
Pink stain: of teak, associated with Fusarium solani, 628
Pinus spp.: rust galls on shortleaf-slash pine hybrids in Texas, similar to but not identical to Cronartium quercuum or C. fusiforme, 248; additional information on pathogenicity of Scleroderris lagerbergii, 738; Diplodia tip blight, occurrence and control in Central Great Plains, 359; Scleroderris canker, Scotch pines less susceptible than red or jack pines, 37
--- albicaulis (whitebark pine): possibly a new host in California for white pine needle cast, Bifusella linearis, 305
--- balfouriana (foxtail pine): white pine blister rust found on, in northwestern California, 391
--- banksiana (jack pine): antibiotic treatments to prevent stem rusts in seedlings, 538; infection of one tree with both eastern and western gall rust, first report of both on same tree, 328
--- clausa (sand pine): susceptible to Fomes annosus in Georgia, 635
--- contorta (lodgepole pine): Lentinellus montanus causes butt rot of heartwood, 820
--- edulis (piñon pine): plant-parasitic nematodes associated with, 52
--- elliottii (slash pine): fusiform rust control on southern pines in nurseries, 597
--- flexilis (limber pine): white pine needle cast in California, possibly new host, 305
--- monticola (western white pine): blister rust control with cycloheximide and phytoactin ineffective, 6
--- ponderosa (western yellow pine): plant-parasitic nematodes in marginal Pinus ponderosa stands in central New Mexico, 52; determining the age of comandra rust infection on, in California, 305; decline and death in southern California apparently due to oxidants, mainly ozone, 707; sodium-linked disease in Denver, Colorado, 908
--- --- var. scopulorum (Rocky Mountain yellow pine): occurrence of lodgepole pine dwarf mistletoe on, in northern Colorado, 125
--- sylvestris (Scots pine): natural infection by lodgepole pine dwarf mistletoe in Canada for first time, 409
Piricularia oryzae: 703
Plantain (see Musa): cooking banana, 901
Plasmopara viticola: 372
Plum (see also Prunus): 'Santa Rosa' plum apparently newly recorded host for European mistletoe, Viscum album, in California, 198; Šarka (plum pox) disease, 250; 253
Podosphaera leucotricha: 754
Pod rot: of cacao, caused by Phytophthora palmivora, 790
Polianthes tuberosa: foliar disease on tuberose in Hawaii caused by Aphelenchoides besseyi, believed to be first report of a foliar nematode on tuberose, 56
Pome spotted anthracnose (see Anthracnose)
Poplar (see Populus)
Populus spp.: reactions to infections by Septoria musiva and Diplodia tumefaciens and to frost injury in Manitoba and Saskatchewan, 829
--- tremuloides: Godronia fuliginosa pathogenic to trembling aspen and willow, 167; Hypoxylon cankers developed on trembling aspen after inoculation, 659
Poria weirii: 961
Postharvest decay: of apples: from Penicillium and Botrytis rot after water treatment, 865; Penicillium rot of McIntosh apples controlled with thiabendazole, 867
---: cantaloup, stem-end rot, control with hot water treatment and chemical fungicides, 262
---: of citrus: Fungicide 1991, thiabendazole and sodium orthophenylphenate compared for control of Penicillium molds in citrus, 623; of Valencia

oranges, antifungal compounds tested for effectiveness in control of green mold, 777; preharvest application of fungicides for control of postharvest decay, 844
Potato: golden nematode: Heterodera rostochiensis detected in western New York, 184; in Panama, 587; key to Heterodera "groups" for aid in identification of golden nematode, 827
---: pink rot, caused by Phytophthora cryptogea, recorded for the first time in Western Australia, in 1966, 280
---: ring rot, rapid serodiagnostic test for the detection of water-soluble phytotoxic glycopeptide produced by Corynebacterium sepedonicum in plant tissues, 502
---: root-knot nematode, pathogenic to 18 varieties tested, 956
Powdery mildew: Fungicide 1991 shows great promise for use against a range of powdery mildews, 754
--- ---: of barley, interaction of 60 varieties of barley with 12 cultures of Erysiphe graminis hordei from four countries, 463
--- ---: of cucurbits, control with Fungicide 1991 by soil drenches and seed treatment, 630
--- ---: of papaya, caused by Oidium caricae, controlled with fungicide + spreader sticker, 814
Pratylenchus penetrans: 57; 383; 519
Preharvest spray: for control of postharvest decay of citrus, 844
Prince Edward Island: 213; 393
Prune (see also Prunus): stem painting with systemic-chemicals on 'Italian' prune suppressed Cytospora infection, 154
Prunus spp. (see also common names): Fungicide 1991 found outstanding in field control of brown rot on stone fruits in California, 722; responses to postplant and preplant soil fumigation in California peach, walnut and prune orchards, 890
---: stem pitting of peach: apparently not previously described, 91; further observations give first report of stem pitting in seedlings of peach, apricot and nectarine, 287; in Pennsylvania, 292; found in one orchard in South Carolina, 907
--- virginiana: mosaic virus of apple, bud transmission of cherry ringspot virus from naturally infected P. virginiana to apple, 794
---: viruses separation of Lambert mottle virus from Prunus ringspot virus in Prunus serotina, in Montana, 184; new virus distinct from any other stone fruit or rose virus mechanically transmissible from rose to stone fruit with symptomless infection, 499; šarka (plum pox) disease, 253; stone fruit viruses occurring in United States compared with šarka (plum pox) disease indigenous to Europe, 250
Pseudomonas aptata: 715
--- phaseolicola: 295; 534
--- solanacearum: 330; 901
Pseudotsuga menziesii: first North American report that Rhizina undulata is pathogenic on Douglas-fir seedlings, 579; Verticicladiella wagenerii found on Douglas-fir in El Dorado Co., California for first time, only second time on this tree species, 393
Psilocybe agrariella var. vaccinii: 87
Psylla pyricola: 363; 924
Puccinia carthami: 876
--- coronata avenae: 59; 205; 235
--- graminis f. sp. avenae: 512; 642; 897
--- --- f. sp. secalis: 512; 642; 897
--- --- f. sp. tritici: 512; 642; 897
--- polysora: 136
--- recondita tritici: 186; 437
Pulse crops (see Legumes)
Purple stain: caused by Cercospora kikuchii on soybean, plowing under soybean debris would eliminate spring inoculum, 931
Pyrenophora teres: 215; 395
Pythium spp.: 72; 209; 334; 577
--- aristosporum: 705
--- myriotylum: 179; 416
--- ultimum: 806

Quebec: 167; 403; 738

Races: apple scab, race 5 of Venturia inaequalis identified in England, 799
---: bacterial blight of cotton, survey in Missouri showed races 1, 3, 4, 11, 12, and 15 present in 1966, 739
---: of common bunt, L-16 designation for new pathogenic race of Tilletia foetida, 569
---: Fusarium oxysporum f. sp. lycopersici, first report of incidence of race 2 infecting tomato in New Jersey, 553
---: of Puccinia graminis f. sp. tritici and P. graminis f. sp. avenae in the United States in 1965, 512; in 1966, 897; in 1967, 642; of P. polysora differentiated in Nigeria and designated NPP. 1 and NPP. 2, 136
---: studies on stabilizing selection in Trichometasphaeria turcica (Helminthosporium turcicum), 427

Radiation: exposure of dormant buds of sweet cherry seedlings with crinkle to gamma radiation, 268
Radopholus similis: 528
Raspberry: crown gall, interaction of Meloidogyne hapla and Agrobacterium tumefaciens in relation to infection of raspberry cultivars, 492; tomato ringspot virus indicated as cause of crumbly berry of red raspberry, 386
Redbud tree (Cercis canadensis): dieback due to automobile exhaust fumes, 744
Resistance: Capsicum, some species show resistance to one or more of three viruses (tobacco mosaic, tobacco etch and potato Y), 381; in corn, to Helminthosporium maydis, apparently new type of resistance observed in Nigeria, 134; in cucurbits, scattered colonies of powdery mildew fungus showed resistance to Fungicide 1991, 630; in lima bean, to Phytophthora phaseoli, lines G 1 and G 2 completely immune to strains A and B, 563
Rhizina undulata: 579
Rhizoctonia sp.: 565
--- solani: 284; 399; 486; 613; 781
Rhizopus sp.: 695
Rhizosphaera kalkhoffii: first report on native blue spruce stands in Arizona, 804
Rhodesia: 99
Rice: bacterial blight, caused by Xanthomonas oryzae, symptomatology, 663
---: blast disease, new medium developed to promote sporulation of Piricularia oryzae, 703
---: Meloidogyne graminicola new pest of rice, common name proposed of rice root-knot nematode, 423
---: sheath blight, seven strains of Pellicularia sasakii with physiological specialization isolated from rice, 323
---: stem rot in India, method for calculating assessment of losses, 963
---: sugarcane mosaic virus, cultivated rice varieties show wide range in susceptibility to strains A, B, D, and H of virus, 775
---: tungro virus disease: in Indonesia as indicated by symptoms, virus-vector interaction and varietal reactions, 122; in India, identification and characterization, 871
Root decay: of red clover, clover root curculios when present with Fusarium spp. increased decay in red clover roots, 213
--- diseases: of pecan, nematode and Pythium spp. associated with feeder root necrosis, 334; Verticicladiella wagenerii pathogenic to Douglas-fir in California, 393
Rosa canina: parasitic nematodes associated with in the United Arab Republic, 149
Rot: black, of sweetpotato, hot-water sprout treatment for the control of scurf and black rot, 475
---: boll rot of cotton: Erwinia aroideae caused severe rotting when inoculated into immature bolls, 14; fungi associated with and their frequency in the Mississippi delta, 582; occurrence of fiber containing fluorescent spots associated with Aspergillus flavus boll rot in the U.S. cotton crop, 1967, 671; parasitic Phytophthora boll rot found, 780; technique for evaluating systemic fungicides and insecticides for control, 835
---: brown stem, of soybeans in North Carolina, 895
---: buckeye, of tomato, occurrence of, and Rhizoctonia soil rot on tomato fruit of 21 varieties or breeding lines, 284
---: Candida, of lemons, not previously described but probably new only in recognition, 433
---: crown, of Peperomia, Phytophthora nicotianiae parasitica causal agent, found in commercial greenhouses in Ohio, 244
---: fruit: of stone fruits, Fungicide 1991 more effective than other fungicides in control of brown rot blossom blight, 722; of strawberry, new fruit rot found in Israel, caused by Pestalotia longisetula, 472; of tomato, control by oils, 555
---: Fusarium stem rot of carnation cuttings, losses from, 840
---: onion white rot, DCNA applied in transplanting furrow appears to give good control in Washington, 272; isolation in pure cultures, 277
---: Penicillium and Botrytis, of apples after commercial water dip before storage, 865
---: Phytophthora, of soybean, response of resistant and susceptible varieties to 2,4-DB herbicide, 355
---: ring rot of potato, caused by Corynebacterium sepedonicum, rapid and specific method for detection of tissues infected with the ring-rot organism, 502
---: root: of alfalfa, Fusarium solani infection related to nutritional stress, 939; of coffee, new record, 773; of

corn, first report of Helminthosporium pedicellatum causing root rot of corn in India, 477; of cotton, comparative fungicidal effectiveness of chemicals on Phymatotrichum omnivorum in laboratory soil cultures, 232
---: ---: of pea: caused by Aphanomyces euteiches, an attempt to associate plant-parasitic nematodes with pea root rot potential, 190; germination of oospores of A. euteiches affected by hydrogen-ion concentration, 264; Fusarium root rot reduced with 1-chloro-2-nitropropane soil-incorporation or in-furrow treatments, 72; Fusarium and Rhizoctonia rot of garden pea controlled with a soil fungicide, 565
---: ---: Pythium root rot pathogenic to Indian ornamental tree, Delonix regia, 705
---: ---: Rhizoctonia root rot of soybeans reached epidemic proportions in Iowa in 1967, 613; of snap bean, greenhouse studies on resistance, 486
---: ---: and stem, pathogenicity of Pythium myriotylum to several grass and vegetable crops, 179
---: ---: stem or fruit, caused by Macrophomina phaseoli, new weed hosts found, 223
---: ---: Texas root rot, most important fungus disease of grapevines in Mexico, 372
---: ---: yellows-root, of spinach, possibly nutritional disorder, 801
---: Sclerotinia flower rot of chrysanthemum in Louisiana, 120
---: stalk: of corn caused by Diplodia maydis, CO_2-powered BB pistol tested for rapid inoculation, 611; of sorghum, resistance or susceptibility of sorghum varieties to red stalk rot pathogen measured by pattern of cell death in stalk tissue, 68
---: stem: of carnation, caused by Armillaria mellea, 415; of rice, caused by Sclerotium oryzae, formulae for determining percentage disease incidence and corresponding percentage loss in yield, 963; of snapbean, in southern Georgia caused by Pythium myriotylum, 416
Rotylenchulus reniformis: 127
Rubbertree: patch canker in Malaysia caused by Phytophthora palmivora, 132
Rumex brownii: found as new host for Verticillium nigrescens, 904
Rust: cyclone collector for collecting rust urediospores modified to facilitate rapid handling of cultures and to improve efficiency of spore deposition, 148; possibly new pine-oak rust in Texas, 248
---: from barberry, physiological races of Puccinia graminis tritici, P. graminis avenae, and P. graminis secalis identified in United States in 1965, 512; 1966; 897; 1967, 642
---: blister: comandra blister rust, on ponderosa pine in California, determining age of, 305
---: ---: white pine: control with antibiotics ineffective, 6; found for first time on foxtail pine in California, 391
---: cedar-apple, of Juniperus scopulorum and J. virginianae, treated with several fungicides for control, 692
---: crown: of oats, Iowa tests indicate additional sources of tolerance to crown rust, 59; nickel-maneb mixture superior to zineb or maneb for control in Iowa, 205
---: ---: of tall oatgrass, good to excellent resistance found in 16 selections among 14 plant introductions and also winterhardiness, 235
---: fir needle rust caused by Hyalopsora aspidiotus, occurrence in western United States and Canada, 336
---: foliage rust of safflower, effect on yield, test weight and percentage of seed oil, 876
---: fusiform: control of on slash pine seedlings in nurseries with ferbam, 597; Cronartium fusiforme infection of slash pine, relation of potential rust infection to high pressure weather systems, 419
---: gall: eastern (Cronartium ribicola): antibiotic treatments not effective on jack pine seedlings in Minnesota, 538; eastern and western gall rust on single jack pine tree in Minnesota, 328
---: ---: western (Peridermium harknessii): of Cronartium coleosporioides complex confirmed in the Maritime Provinces, 455; identified on Pinus banksiana from Quebec and New Brunswick, 650
---: leaf: of corn, two physiologic races of Puccinia polysora found in Nigeria, 136; of wheat, method for rapid classification of cultures and method of data processing, 437
---: ---: and stem of wheat, control with DCMOD in combination with foliage sprays, 856
---: stem: of oat and wheat, physiological races of Puccinia graminis avenae and Puccinia graminis tritici in the United States in 1965, 512; 1966, 897; 1967 642
---: sweetfern rust, Cronartium comptoniae, 538

Safflower: Puccinia carthami infection on lower leaves only, did not significantly reduce yield, test weight or oil percentage in Utah tests, 876
Salt damage: to vegetation in Washington, D. C. area from winter-salting of roadways, 350
Sarka (plum pox) disease: comparison to stone fruit viruses occurring in the United States, 250; most serious virus disease of European plum, 253
Saskatchewan: 829
Saussurea candicans: leafspot caused by Alternaria tenuis on, 248
Scab: apple: decomposition of fall leaves by nitrogenous fertilizers reduces production of ascospores from apple leaves in spring in Connecticut, 728; Fungicide 1991 effective in inhibiting perithecial development of Venturia inaequalis after single fall application, 654; identification of race 5 of Venturia inaequalis, 799
---: of pecan: new fungicides give good control in Georgia, 108; 144 varieties of pecan screened for resistance to scab and bunch disease, 307
Scleroderris lagerbergii: 37; 738
Sclerophthora rayssiae zeae: 446
Sclerospora macrospora: 569
--- sacchari: 446
Sclerotinia homoeocarpa: 410
--- sclerotiorum: 120
--- trifoliorum: 882
Sclerotium cepivorum: 272; 277
--- oryzae: 963
Scurf: of sweetpotato, treatment of basal portions of sweetpotato sprouts grown from roots for 2 to 5 minutes in water at 120°F completely eliminated scurf fungus, 475
Scutellonema brachyurum: first known report as a parasite on soybeans, 40
Seed: effect of soil fungicides on development of fungi on kernels of peanut, 718; oil from safflower, oil percentage of seed harvested from rusted plots tends to be slightly higher than from unrusted plots, 876
---: transmission: of maize dwarf mosaic virus in corn, found to be very low, 863; of peanut stunt virus, selected symptomless plants showed low incidence, 522
---: treatment: anaerobic, of barley seed to kill loose smut fungus, resulted in lower germination after storage, 792; peanut seed treatment with fungicides, effect on seed mycoflora, germination and seedling emergence, 240; of squash and cucumber with systemic fungicide for powdery mildew control, 630
Seedborne: bacterial blight of cotton, acid-delinting and fumigation not completely effective in eradication of pathogen, 981; mimosa wilt can be seedborne, only known record of wilt-inducing Fusarium being found within seed of woody plant, 949; Verticillium dahliae seed infection of Xanthium pungens in New South Wales, 976
Seedling: blight of corn: factors affecting severity, 79; fungi associated with in United Arab Republic, 84
---: reaction of barley, oats and wheat to Pythium species, 209; cotton soreshin caused by Rhizoctonia solani, fungicides tested for chemotherapeutic activity, 399
Septoria apiicola: 142
--- arabidis: 175
--- musiva: 829
--- sisymbrii: 175
Serum: anti: of tobacco mosaic virus produced in horse in California, 860; use of antiserum for detection of ring rot infection in potato tissues, 502
Sesamum orientale: phyllody and leaf curl viruses of, control with insecticides, 367
Sesbania exaltata: as summer cover crop, 573
Shuck disease: of pecan, appears to be result of physiological-nutritional imbalance, not an infectious disease, 189
Sitona hispidula: effects of clover root curculios and Fusarium species on red clover, 213
Smut: common corn smut, Ustilago maydis caused unusual epiphytotic condition in Indiana and Illinois, 922
---: flag smut, Urocystis agropyri, and stripe smut, Ustilago striiformis, found together on 'Merion' Kentucky bluegrass in Ohio, 633
---: head smut of corn, chemical control of Sphacelotheca reiliana with in-furrow treatment, 726
---: stripe, together with flag smut on 'Merion' Kentucky bluegrass turf, 633; Fungicide 1991 effective in control of, 635
Sodium: foliar chlorosis and tipburn of ponderosa and other pines, high level of sodium found in diseased foliage possible cause, 908
Soil: low soil moisture and the predisposition of tobacco plants to blue mold infection, 310
---: amendments: alfalfa "green" manure and steer manure reduced root-knot

disease in outdoor microplots during 4-year period, 315; white pine sawdust, paper and cottonseed meal reduced larval invasion (Heterodera tabacum) of eggplant roots, 441; petroleum mulch helped reduce damping-off of cucumber, 806
Soil: treatments: application of granular organophosphate and carbamate pesticides controlled sting and stubby-root nematodes on vegetables, 19; broad spectrum fumigants and mixtures of specific pesticides tested for possible multiple pest control, 193; for control of reniform nematodes in cotton, several nematocides tested, 786; after harvest plowing would remove source of spring inoculum of Cercospora kikuchii on soybean, 931
---: ---: 1-chloro-2-nitropropane (NIA 5961) + PCNB as in-furrow application gave good control of head smut of corn, 726; soil incorporation and in-furrow treatments for control of pea root rot, 72
---: ---: DCMOD, in-furrow application at planting for control of early infection by leaf and stem rust of wheat, 856
---: ---: DCNA applied in transplanting furrow gave good control on onion white rot, 272
---: ---: disulfoton, phorate and carbamate, UC 21149 granular formations applied with seed or as side dressing, ineffective in control of Verticillium wilt, 281
---: ---: Fungicide 1991: applied in-furrow reduced root rot in garden peas and increased yield, 565; as soil drench to control powdery mildew of cucurbits, 630; gave protection against postemergence damping-off of cotton seedlings in greenhouse tests, 781
---: ---: methyl bromide-chloropicrin-propargyl bromide (Trizone) appeared to give some control of crown gall of mazzard cherry seedlings, 102; methyl bromide and chloropicrin (MBR-CP) reduced soilborne wheat mosaic virus incidence in 'Pawnee' wheat in Kansas, method might be useful where wheat is cultured intensively in small plots, 559
Solanum melongena (see also Eggplant): little leaf disease in India, method for estimating severity of disease, 118
--- nigrum: black nightshade new host for Verticillium nigrescens in Australia, 904
Sooty mold: of cotton and other hosts, disease not previously described from India, 602
Sorghum: anthracnose, reaction of selected sorghum lines to artificial inoculation of Colletotrichum graminicola, 164
---: maize dwarf mosaic virus, first identification in New Mexico, 365
---: plant-parasitic nematodes associated with, in fields of great millet in India, 396
---: red stalk rot, pattern of cell death in stalk tissue good predictive measurement for resistance or susceptibility to spread of pathogen, 68
South Carolina: 486; 571; 907; 926
South Dakota: 349; 795
Soybean: "Bibliography of Soybean Diseases" now available, 256
---: brown stem rot incidence in North Carolina, 895
---: cotyledon necrosis, disease apparently caused by weathering, 459
---: nematode: associations, distribution and occurrence in southeastern U.S.A., 40; Hoplolaimus columbus found to cause severe stunting in South Carolina, 571
---: Phytophthora rot, resistant varieties appear to possess resistance to damage from 2,4-DB (4 (2,4-dichlorophen=oxy)butyric acid, 355
---: purple stain, survival of Cercospora kikuchii on stems in field, 931
---: Rhizoctonia root rot in Iowa of epidemic proportion, 613
---: soybean mosaic and bean pod mottle viruses, effect of single and double infections, 344
Spelling change: change of spelling from "nematocide" to "nematicide", 984
Sphacelotheca reiliana: 726
Spinach: viruses of, in Texas, 761; yellows-root rot, in Connecticut, cause not yet known, 801
Spongospora subterranea f. sp. nasturtii: 859
Spores: collecting fungus spores with cyclone collector, 148
Spruce (see Picea)
Squash (see Cucurbits)
Stalk rot: of corn, Diplodia stalk rot inoculating technique for corn, 611
Stem pitting: of peach, characterized by foliar symptoms of girdling, lower trunk enlargement and extremely thickened bark and wood pitting, 91; seedlings of peach, apricot and nectarine found affected, first record of occurrence in seedlings, 287; in Pennsylvania orchards, cause of disease

yet unknown, 292; found in 'Coronet' peach trees in Spartanburg County, South Carolina, 907
Stem tip cultures: use of apical meristem and callus cultures to produce geranium plants, 600
Stereum sanguinolentum: 489
--- taxodii: 482
Stone fruits: brown rot, efficacy of Fungicide 1991 in reducing fruit rot, 722
Storage: soybean cyst nematodes survived in seed bags after 17, 18, and 22 months, 45, cryogenic storage of Phytophthora phaseoli, 507; barley seeds treated anaerobically suffered germination loss more rapidly than untreated seeds, 792
---: decay: effect of ozone on Penicillium mold decay and sporulation, 245; control of gray-mold of Barlinka grapes with vapor-phase chemical 2-acetyl-3-hydroxyfuran (GS-7174), 479; control of ripe fruit rots in tomato with oil-soap emulsion sprays, 555; market deterioration of cultivated mushroom and microorganisms associated with, 712; postharvest decay of Florida citrus controlled with Fungicide 1991, 844; thiabendazole as postharvest fungicide controlled Penicillium rot of apple, 867
Strains: physiological specialization of seven strains of Pellicularia sasakii, 323; development of two new lines of lima bean, G 1 and G 2, completely immune to strains A and B of Phytophthora phaseoli, 563; sugarcane mosaic virus, 1:3 and 1:10 dilutions of inoculum for screening seedlings for resistance to strain H may be more efficient, 705; sugarcane mosaic virus, A, B, D, and H, reaction of different rice varieties to, 775; latent strain of tobacco mosaic virus, 919
Strawberry: fruit rot, caused by Pestalotia longi-setula, apparently first record of pathogen on strawberry fruit, in Israel, 472
---: viruses: transmission of strawberry mottle virus by juice and aphids to herbaceous hosts, 64; virus infection and its effect on the strawberry cultivar Northwest, 811
Streptomyces griseus: 3
Stripe disease: of winter wheat, caused by Cephalosporium gramineum, in Washington, 590; distribution of soilborne disease of wheat in Montana, 818
Stripe smut (see Smut): 633; 635
Sudan: 405
Sugarbeet: downy mildew, disease of minor importance in California since the introduction of more resistant varieties, 297; Longidorus africanus increased more on sugarbeet than other crops in 5-month test, 748; study of sugarbeet virus complex in southwestern Idaho, 965
Sugarcane: cyst nematode, Heterodera sacchari, causes stunting of plants in Nigeria, 237; first record of cyst nematode, on sugarcane in India, 626
---: leaf scald, caused by Xanthomonas albilineans found in Florida for first time, 646
---: mosaic virus: strain I found in Louisiana, 449; inoculum concentration in relation to screening seedlings for levels of resistance to strain H, 705
Surface sterilization: mycoflora of peanut kernels after surface-sterilization of pods or kernels with mercuric chloride, 0.1%, 608
Surfactant: Tween 20 added to thiabendazole or Fungicide 1991 increased effectiveness against Verticillium in cotton, 944
Survey: aerial survey of eastern Nebraska and Kansas for oak wilt, 357; distribution and occurrence of nematodes associated with soybeans in Georgia, Florida and Alabama, 40; of plant-parasitic nematodes associated with roots of great millet in India, 396; for presence of peanut stunt virus in South Carolina in 1967, 926
Survival (see also Longevity): susceptibility of parasitic nematodes to subfreezing temperatures, Heterodera tabacum most cold-resistant species tested, 768
Sweetfern: rust, caused by Cronartium comptoniae, 538
Sweetgum: bleeding necrosis, Botryosphaeria ribis, as causal agent confirmed, 223; fumigation of nursery beds with methyl bromide improved sweetgum growth by reducing population of soil-inhabiting pathogens, 483
Sweetpotato: several nematocides tested for control of reniform nematodes, 127
---: scurf and black rot, hot-water sprout treatment a promising control for, 475
---: russet crack virus, natural transmission under exposed conditions, 967
Sycamore: pathogenicity of sting nematodes to sycamore seedlings artificially inoculated, 523

Taiwan: 323
Take-all: of wheat, occurrence of in the Pacific Northwest, 716
Tanzania: 628
Taxodium distichum (see Baldcypress)
Taxus canadensis: Phacidium taxicolum on foliage and bark of dead twigs and branches of Canadian yew, in Quebec, 403
Tea (see Camellia sinensis)
Teak: Fusarium solani consistently isolated from collar and stem cankers of mature teak in Tanzania, 628
Temperature: viability of urediospores of Puccinia recondita on dry wheat foliage affected by temperature, 186; relation to development of southern corn leaf blight on sweet corn ears in transit, 847; relation of low temperature to unusual epiphytotic of common corn smut in Indiana and Illinois, 922; response of 'Bel W3' tobacco is an inverse relation between sensitivity to phytotoxicant ozone and temperature in experimental chamber, 957
---: hot water treatments: hot water reduced stem-end rot of cantaloup, 262; of sweetpotato sprouts, promising control for scurf and black rot, 475
---: subfreezing: Phytophthora phaseoli kept in viable and pathogenic state for 476 days at -60°C, 507; effect on parasitic nematodes, 768
Tephrosia vogelii: a potential source of rotenone found susceptible to several common plant viruses, 530
Texas: 164; 248; 375; 761
Tilletia foetida: 569
Tipburn: or chronic foliar necrosis in Pinus ponderosa in Denver, Colorado, cause unknown, 908
Tobacco: diseases of Florida flue-cured tobacco in 1968, 984; diseases of in Kentucky, 1967, 589; response of 'Bel W3' to ozone in plant fumigation chamber, 957
---: blue mold: occurrence in burley tobacco-growing areas in Kentucky in 1967, 204; plants cultivated under low soil moisture regime more predisposed to infection, 310
---: brown spot, first occurrence in certain dark and burley tobacco-growing areas of Kentucky in 1966, 71
---: viruses: potato virus Y caused veinbanding in Florida sun-grown plants, 887; effect of dimethyl sulfoxide on tobacco mosaic, 61; tobacco mosaic virus, competitive characteristics of latent strain in relation to common TMV, 919
Tomato: diseases of in Colorado in 1967, 542; ripe fruit rots and their control by oils in India, 555
---: anthracnose and early blight, reduction of in Delaware after fungicide treatments, 685
---: bacterial canker: found in Mendoza province, Argentina for first time, 595; seedlings can become infected with as few as five cells of the organism when introduced directly into xylem tissue, 741
---: bacterial spot, use of commercial implementation for successful control, 339
---: bacterial wilt, spread of disease from diseased to healthy plants by clipping, 330
---: buckeye rot, observational differences in susceptibility of tomato varieties to Rhizoctonia rot, 284
---: Fusarium oxysporum f. sp. lycopersici race 2 in New Jersey, apparently first report of race 2 of the fungus in northeastern tomato-growing region, 553
---: Phoma leaf spot, occurrence of foliar infection on tomato and potato in Hawaii, 946
---: root knot, weed and fungus control, evaluation of mixtures of specific pesticides and broad-spectrum fumigants, 193
---: viruses: curly top virus, restricted local spread in Washington, 17; enation leaf curl designated Nicotiana virus 10A or enation leaf curl virus of tomato, 595; several on tomato in Argentina, 674
---: tomato bushy stunt virus found in Argentina, 676
Toxicity: of boron to chrysanthemums due to boron content of water supply, 824
Transmission: maize dwarf mosaic virus seed transmission, in corn, 863; peanut stunt virus, transmission through selected symptomless plants very low, 522; russet crack virus, of sweetpotato, 967; strawberry green petal virus in the field at Prince Edward Island, 393; of Streptomyces phage by saprozoic nematodes, 3; tomato bushy stunt virus, unsuccessful with Olpidium, 379
---: mechanical: of exocortis virus of citrus by contaminated tools, 935; onion yellow dwarf virus, contaminated knife can transmit virus from onion to onion, 581; rose virus capable of symptomless infection of stone fruits, 499; strawberry mottle virus, isolates

of SMV transmitted mechanically or by aphid vector Chaetosiphon jacobi, 64
Tree tomato (Cyphomandra betacea): control of powdery mildew with systemic fungicide, 754
Trichodorus spp.: 149
--- christiei: 19; 573
Trichometasphaeria turcica (Helminthosporium turcicum): 427
Tsuga heterophylla (western hemlock): occurrence of Fomes annosus in intensively managed young-growth stands in western Washington, 370
Tuberose (see Polianthes tuberosa)
Turnip (see Crucifers)
Tylenchorhynchus spp.: 149
--- claytoni: 57
Tylenchus exiguus: 52

Ultraviolet: as a sampling aid for detection of bean seed infected with Pseudomonas phaseolicola, 534
United Arab Republic: 79; 84; 149; 150
Urocystis agropyri: 633
Ustilago maydis: 922
--- nuda: 792
--- striiformis: 633; 635
Utah: 876

Vectors associated with: curly top virus in tomatoes, leafhopper Circulifer tenellus, 17; enation leaf curl of tomato, transmitted by sweetpotato whitefly, Bemisia tabaci, 595; pear decline virus, Psylla pyricola, 363; phyllody and leaf curl viruses of Sesamum orientale, Deltocephalus sp. and Bemisia tabaci, 367; squash mosaic virus, in Venezuela, beetle Acalyma thiemei thiemei found to be new vector of, 171; strawberry green petal virus, Macrosteles fascifrons and Aphrodes bicinctus, 393; strawberry mottle virus, aphid Chaetosiphon jacobi, 64
Veinbanding: caused by potato virus Y found on tobacco in north Florida, 887
Venezuela: 171
Venturia inaequalis: 654; 728; 799
Vernonia anthelmintica: several common plant viruses pathogenic to potential seed oil plant, 530
Verticicladiella wagenerii: 393
Verticillium albo-atrum: 281; 944
--- dahliae: 910; 976
--- nigrescens: 904
Virginia: 162; 914
Virus: four potential new crop species for U.S. agriculture found susceptible to common plant viruses, 530
---: Agropyron mosaic, circular or tubular inclusions detected with electron microscope may help differentiate this virus from wheat streak mosaic virus, 139
---: of apple, mosaic virus of apple, Prunus virginiana a natural host for, 794
---: bean pod mottle virus on soybean, 344
---: bean yellow mosaic virus on cowpeas, 34; 509
---: Bidens mottle virus, from electron microscopy virus appears to be previously undescribed and to belong to the potato virus Y group, 763
---: crumbly berry of raspberry, tomato ringspot virus suggested as cause, 386
---: exocortis virus: found in Ortanique orange in rootstock trial in Jamaica, 741; of citrus in India, 834; survived 8 days on steel knife blade, 935
---: corn: maize dwarf mosaic virus: air-blast inoculation of corn varieties with virus in disease nursery affords high level of infection, 24; use of corundum abrasive essential with mobile air-blast inoculator for high level of infection, 28; presence of virus in seven counties in Georgia established, 320; on corn, broomcorn, sorghum and Sudan grass, first report of occurrence in New Mexico, 1967, 365; dent corn synthetics resistant to mosaic viruses and other corn diseases in Pennsylvania selections, 424; two strains designated A and B recovered from dent and sweet corn in New Jersey, 496; Pennsylvania studies on interaction of virus-host-pesticides under field conditions, 545; seed transmission, 863
---: of cowpeas: cucumber mosaic virus, cowpea chlorotic mottle virus and bean yellow mosaic virus in Alabama, 34; bean yellow mosaic virus, cowpea chlorotic mottle virus, and cucumber mosaic virus tested for effect upon growth and seed production of 'Clay' and 'Early Ramshorn' cultivars, 509
---: cucumber mosaic on cowpeas, 34; 509
---: of cucurbits: cucurbit virus complex in Kansas seems to differ from cucurbit complexes reported from other U.S.A. sections, 11; viruses occurring in Venezuela, 171
---: of grape, leafroll, fan leaf and corky bark of grapevines in Mexico, 372
---: oat mosaic, influence of crop rotation and oat mosaic virus on yield and test weight of oats in South Carolina, 430

---: onion yellow dwarf, found to be mechanically transmissible to mother bulbs by use of contaminated knife, 581
---: of pea, disease reaction of Pisum sativum plant introductions to alfalfa mosaic virus (AMV), red clover vein-mosaic virus (RCVMV) and Wisconsin pea streak virus (WPSV), 160
---: of peanut, "bud necrosis" suggested as name for possibly new virus, in India, 494
---: ---: mottle, no immunity found in 465 introductions in Georgia tests, but some differences in resistance indicated, 467
---: ---: stunt virus: transmission of virus through seed from selected symptomless peanuts, 522; none of 300 varieties tested showed immunity but some differential response indicated, 914; virus-free peanut fields in South Carolina despite PSV in white clover throughout most of State, 926
---: pear decline, leaf drop a new symptom seen in own-rooted 'Magness' pear inoculated with pear decline virus, 363
---: pear leaf curl, evidence suggests pear leaf curl and pear decline may be caused by same virus, 924
---: of pepper: in Mendoza, Argentina, 674; pepper varietal reaction to naturally occurring viruses in California, 928
---: Prunus: cherry: cherry rasp-leaf virus, in California, 618; cherry ringspot virus, agent for mosaic virus of apple with Prunus virginiana a natural host, 794; Lambert mottle virus separated from Prunus ringspot virus on Prunus serotina (choke cherry), 184; necrotic ringspot of sour cherry, record of increase and spread in orchard in Pennsylvania, 972
---: ---: prune: Šarka (plum pox) disease in reference to virus diseases of Prunus in the United States, 250; 253
---: potato virus Y: some South American Capsicum species show resistance to, 381; Bidens mottle virus appears to belong to potato virus Y group, 763; on tobacco in north Florida, 887
---: of rice: tungro disease of rice in Indonesia, 122; in India, 871
---: of rose, new virus causes green mottled foliage on Rosa rugosa, distinct from any other stone fruit or rose virus, 499
---: of Sesamum orientale, phyllody and and leaf curl control with insecticides, 367
---: soybean mosaic and bean pod mottle viruses, effect of single and double infections on yield and seed characters, 344
---: of spinach, beet curly-top, spinach yellow-dwarf and tobacco ringspot viruses most prevalent on spinach in Texas in 1966-1967, 761
---: of strawberry: mild yellow edge, crinkle, mild yellow edge plus mottle, and mild yellow edge plus crinkle reduced number of runners of strawberry cultivar 'Northwest', 811; strawberry green petal or clover phyllody virus, most critical period of infection in strawberry fields is mid-August, 393; strawberry mottle virus transmitted mechanically, or by aphid vector, Chaetosiphon jacobi, from strawberry to Chenopodium, 64
---: of sugarbeet, an attempt to identify causal entity of yellow beet complex in Idaho, 965
---: sugarcane mosaic: strain I, a new designation in Louisiana, 449; concentration of 1:3 or 1:10 suggested for screening for levels of resistance to strain H, 705; wide range in susceptibility of cultivated rice varieties to four strains of SCMV in Louisiana, 775
---: of sweetpotato, russet crack, 967
---: tobacco mosaic: number of lesions induced by virus affected by varying concentrations of dimethyl sulfoxide (DMSO) in inoculum, 61; TMV, tobacco etch and potato Y viruses, some South American Capsicum species show resistance to, 381; water extracts from most fungi tested (mostly mushrooms) contained substances that inhibit TMV, 528; production of (Anti-TMV) antiserum in a horse, 860; further study of latent strain of TMV, 919
---: tobacco ringspot, pathogenic to Luffa acutangula in host range studies, 374
---: of tomato: curly top virus in tomatoes, its spread during low temperature, 17
---: ---: enation leaf curl considered to be strain of tobacco leaf curl virus, 595
---: ---: bushy stunt: transmission unsuccessful with Olpidium, 379; of tomatoes and peppers in Mendoza, Argentina, first report of occurrence of tomato bushy stunt, 674; first report of occurrence in Argentina, 676
---: tomato ringspot virus on red raspberry, suggested as cause of crumbly

berry, 386
---: turnip mosaic virus, found in Florida, 759
---: of wheat: soilborne wheat mosaic, effect of soil fumigation on incidence and on wheat yield, 559; wheat streak mosaic virus, evaluation of 12 varieties of spring wheat in Montana, 201
Viscum album: European mistletoe rediscovered on apple and on several other host trees in California, 198

Walnut: fumigation studies with preplant and postplant treatments for control of nematodes, 890
Washington: 272; 370; 590; 983
Washington, D. C.: 350; 792
Watercress: crook root, found in commercial watercress beds in Florida and Pennsylvania, 859
Watermelon (see Cucurbits)
Weather: high-pressure systems related to fusiform rust infection in southeastern United States, synoptic and prognostic charts should improve weather forecasts for rust control programs, 419
Western gall rust (see Rust)
West Indies: 741
Wetwood: of cottonwood tree, in lower Mississippi Valley, 822
Wheat: bunt, common, new race of Tilletia foetida found in Idaho, 569
---: Cephalosporium stripe disease: in Washington, factors influencing stripe disease within its area of adaptation, 590; one of most important soilborne diseases in Montana, 818
---: downy mildew, caused by Sclerospora macrospora, first record of in India, 569
---: powdery mildew, controlled with systemic fungicide, 754
---: Pythium aphanidermatum most pathogenic of five Pythium species to barley, oat and wheat seedlings, 209
---: rust: leaf: Puccinia recondita tritici, viability of urediospores on dry wheat foliage affected by temperature, 186; mechanical planting and inoculation of wheat seedlings, also method of data processing, 437
---: ---: ---: and stem, rust control by combinations of DCMOD with foliage sprays, 856
---: ---: stem: Puccinia graminis tritici, annual survey of physiologic races in the United States in 1965, 1966, 1967, 512; 642; 897
---: take-all caused by Ophiobolus graminis increasing in importance in Pacific Northwest with increase of irrigated wheat, 716
---: viruses: soilborne wheat mosaic virus, soil fumigation and its effect upon incidence of SBWMV and wheat yield, 559; wheat streak mosaic virus, yield losses in 12 varieties of spring wheat evaluated, 201
Wilt: bacterial, of tomato, greenhouse studies of spread by clipping, 330
---: Cephalosporium wilt of American elm found in Mississippi and Arkansas, 170
---: Fusarium: of alfalfa, nutritional stress increases susceptibility, 939; of tomato, apparently first report of incidence of race 2 in New Jersey, 553
---: of mimosa: caused by Fusarium oxysporum perniciosum, incidence in Arkansas confirmed, 79; found to be carried in seed, 949
---: oak: caused by Ceratocystis fagacearum, distribution in eastern Kansas and Nebraska, 357; first report in Platte County, Missouri, 811
---: of persimmon, detection of several diseased trees in Arkansas, 79
---: Phytophthora, of hops in California, wilt phase of Phytophthora crown and root rot of hops, 76
---: of cotton: caused by Verticillium albo-atrum, three organic phosphate systemic insecticides had little effect upon, 281; surfactant Tween 20 added to systemic fungicides increased control of fungus, 944
---: --- dahliae: and Fusarium oxysporum f. vasinfectum together on cotton plants increased severity of wilt and reduced incubation period, 910; seeds of Xanthium pungens infected with V. dahliae an important host for fungus in cotton-growing areas in New South Wales, 976
Wisconsin: 160; 341

Xanthium pungens: important host of Verticillium dahliae in cotton-growing areas of New South Wales, 976
Xanthomonas albilineans: 646
--- malvacearum: 739; 981
--- oryzae: 663
--- vesicatoria: 683
Xiphinema spp.: 149
--- americanum: 52; 795
--- elongatum: 150
--- index: 626

Yellows-root rot: of spinach, 801
Yew (see Taxus)
Yugoslavia: 250; 253

CROPS RESEARCH DIVISION

Prepared by Marguerite G. Benedict

☆U. S. GOVERNMENT PRINTING OFFICE : 1969 343-155/12

UNITED STATES DEPARTMENT OF AGRICULTURE
AGRICULTURAL RESEARCH SERVICE
Crops Research Division
Beltsville, Maryland 20705

Postage and Fees Paid
U. S. DEPARTMENT OF AGRICULTURE

Official Business